

Electromagnetic pion and kaon form factors in light-cone resummed perturbative QCD

Udit Raha* and Andreas Aste†

Department of Physics, University of Basel, Switzerland.

(Dated: March 1, 2009)

Abstract

We analyze the electromagnetic pion and kaon form factor by including radiative and higher-twist effects within the framework of resummed pQCD in the space-like region. We focus on the transition from the perturbative to non-perturbative behavior in the phenomenological intermediate energy regime. Using a modified “ k_T ” factorization scheme with transverse degrees of freedom, we evaluate the non-perturbative soft contributions as distinct from the hard contributions, ensuring no double counting via the Ward identity at $Q^2 = 0$. The soft contributions are obtained via local quark-hadron duality while the hard contributions rest on the well known collinear factorization theorem using model wave functions with modified Brodsky-Huang-Lepage type ansatz and distribution amplitudes derived from light-cone QCD sum rules. Our analysis shows that the perturbative hard part prevails for large Q^2 beyond 50-100 GeV², while for low and moderate momentum transfers below 10-16 GeV² the soft contributions dominate over the hard part. Thus, we demonstrate the importance of including the soft contributions for explaining the experimental form-factor data.

PACS numbers: 12.38.Bx, 12.38.Cy, 12.39.St, 13.40.Gp

*Electronic address: Udit.Raha@unibas.ch

†Electronic address: Andreas.Aste@unibas.ch

I. INTRODUCTION

During the past decade, QCD-oriented studies have been shifting steadily toward exclusive channels. For a long time the electromagnetic structure of pions has been subjected to numerous experimental and theoretical investigations through the study of electroproduction reactions. Extensive experimental studies of pion electroproduction reactions like $ep \rightarrow e\pi^+n$ or $en \rightarrow e\pi^-p$ have been carried out in the past at CERN, Cornell, DESY and more recently at JLab Facility [1, 2, 3, 4, 5, 6, 7, 8]. Since the mid-90s, kaon electroproduction reactions like $A(\gamma, K)YB$ and $A(e, e'K)YB$ (A is the target, Y the produced hyperon and B the recoil) have also attracted renewed interest in nuclear physics at both experimental [9, 10] and theoretical [11, 12] level. The main ingredients for the description of electromagnetically induced kaon production are embedded in the so-called Chew, Goldberger, Low and Nambu (CGLN) scattering amplitudes. In case of the longitudinal component of the electron induced unpolarized differential cross-section, the t-channel diagram dominates and (in certain kinematic conditions) can be factorized [9, 13] as $\sigma_L = k \cdot \mathcal{F}(Q^2) \mathcal{G}(W) \mathcal{H}(t)$, where k is a kinematic factor, $\mathcal{F}$, $\mathcal{G}$, and $\mathcal{H}$ are functions of the 4-momentum transfer squared Q^2 of the virtual photon, the invariant mass W and the Mandelstam variable t , respectively. The function $\mathcal{F}(Q^2)$ implicitly contains the information about the electromagnetic form factor of the kaon. Note that $\mathcal{F}(Q^2)$ is not the actual form factor, but rather a complicated function from which the form factor can be extracted using e.g., Chew-Low extrapolation and deconvolution algorithms. A precise knowledge of the form factor is of fundamental importance for a realistic and accurate description of *exclusive reaction* mechanisms and plays a key role in understanding the interplay between perturbative and non-perturbative physics at intermediate energies. Moreover, the study of form factors provides direct insight into the electromagnetic structures and charge distributions of hadrons as they couple with photons.

To date, the electromagnetic kaon form factor is very poorly known and only measured at very low Q^2 (below 0.2 GeV²) [5, 14]. The status for the (quasi-free) Lambda (Λ) and Sigma (Σ) hyperons is even worse, i.e., there are simply no available experimental data. Basic quantities like the strong coupling constants $g_{K\Lambda N}$ and $g_{K\Sigma N}$ derived from purely hadronic processes or theoretical considerations are not well established and must be considered adjustable. Recently, however, there appeared quite large and precise data sets on photo-production of kaons from the SAPHIR (ELSA) [15], CLAS (CEBAF) [16] and

LEPS (SPring8) [17] collaborations. There is also new data on electroproduction of positive kaons from experiment E98-108 at CEBAF which are being analyzed at the moment.

Keeping in mind the increasing accuracy of experimental data, an accurate theoretical description of the electromagnetic form factors of pseudo-scalar charged mesons at intermediate energies is of primal importance. To our knowledge, especially for the kaon there are very few theoretical works in this direction [18, 19, 20, 21]. In this paper, in addition to the pion form factor, we analyze the kaon form factor for a broad range of space-like momentum transfer. Our framework is based on resummed perturbative light-cone QCD formalism [22, 23], unlike conventional approaches like “asymptotic” and lattice QCD or from sum rules that rely on many unchecked hypotheses. The experimental results could then be used to extract the various *distribution amplitudes* (DAs) [24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41] used in the above formulation. Of course, only a handful of experimental hadron electroproduction data points are presently available to make definitive statements on the validity of different theoretical approaches. Since in almost all cases, the corresponding data points are merely concentrated in the very low energy region ($Q^2 < 1 \text{ GeV}^2$), perturbative QCD (pQCD) has limited predictive power due to the rapidly growing magnitude of the strong coupling, as Q^2 tends to zero. Despite the existing plethora of literature on the predictions on electromagnetic meson form factors based on various approaches (see e.g., Refs. [21, 22, 23, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72] for the pion and Refs. [18, 19, 20, 21] for kaon form factors, to give a highly incomplete list of references), till date there is considerable amount of debate as to their exact behavior in the phenomenological low and moderate energies between $Q^2 \approx 4\text{-}50 \text{ GeV}^2$. Nevertheless, we try to give our assessment to the existing scenario and try to explain the experimental data first for the pion form factor, where statistics are far more decent as compared to that of the kaon. Then we extend our analysis to the kaon form factor, where experimental data is still too limited for any meaningful comparison. Hopefully, with the planned 12 GeV upgrade proposal of the CEBAF experiment (at JLab) in the near future, studies of intermediate energy QCD can prove to be fruitful.

The standard “asymptotic” QCD is known to make successful predictions of many phenomena like *dimensional scaling*, *helicities*, *color transparency*, etc. for exclusive processes, as Q^2 tends to infinity [47, 70, 71, 72, 73]. The approach relies on the so-called *collinear*

factorization theorem [71, 72] which provides an outstanding way of isolating the partonic part accessible to pQCD from the non-perturbative parts. The basic ingredients are: (a) the hadron DA ϕ , which encodes the non-perturbative information regarding the momentum distribution of the constituent “near” on-shell valence partons collinear to the hadron and also features of the QCD vacuum structure as expressed through the quark condensates [74, 75, 76], and (b) a scattering kernel T_H , describing the hard scattering of “far” off-shell valence partons. The overall amplitude of the exclusive process is then given by the convolution, $\phi \otimes T_H \otimes \phi$. However, the application of pQCD to exclusive processes at intermediate momentum transfers or phenomenologically accessible energies (e.g., at CEBAF) has been the subject of severe controversies and criticisms [69, 77, 78, 79, 80, 81, 82, 83, 84, 85]. It is widely anticipated that non-perturbative effects arising from soft gluon exchanges or from endpoint contributions to phenomenologically acceptable wave functions (or DAs) dominate and may severely preclude the predictability of pQCD. Hence, in this paper we use a modified “resummed” pQCD formalism (as proposed in Refs. [22, 23]) which is believed to largely enhance the predictability of pQCD in a self-consistent way at intermediate energies. The central issue here is the inclusion of transverse momentum k_T dependence that necessitates the inclusion of a *Sudakov suppression* factor. This is to organize the large double logarithms of the type $\alpha_s \ln^2 k_T$, arising at all orders due to the overlap of soft and collinear contributions of radiative gluon loop corrections. Such a resummation effectively suppresses the non-perturbative contributions at large energies. However, this may not still be effective enough when talking of Q^2 down to a few GeV^2 . These facts are in agreement with some of the recent findings, reported in Refs. [62, 63, 64] for the pion form factor and also in the context of B systems [86].

In this paper, we emphasize the importance of including two distinct contributions to exclusive quantities for obtaining good agreement with experimental data at low and moderate energies: firstly, the non-factorizable soft contributions which are not calculable within the perturbative framework and secondly, the power suppressed corrections from non-leading twist structures (twist-3) determining the preasymptotic behavior. In other words, the electromagnetic form factor $F_M(Q^2)$ for a charged meson M should be written as

$$F_M(Q^2) = F_M^{\text{soft}}(Q^2) + F_M^{\text{hard}}(Q^2), \quad (1)$$

where $F_M^{\text{hard}}(Q^2)$ is the factorizable part computable in pQCD and $F_M^{\text{soft}}(Q^2)$ is the non-

factorizable soft part. The soft contributions to the form factors can be calculated using phenomenological quark models either incorporating transverse structure (momentum) dependence of the hadron wave functions (see e.g., Refs. [60, 78, 80, 81, 87]) or from QCD sum rules via *Local Duality* (see e.g., Refs. [48, 49, 50, 51, 61, 88].) In the present paper, we follow the latter approach. We also focus on the presence of non-perturbative enhancements arising from kinematic endpoint regions of the scattering kernel which tend to invalidate collinear factorization. For our calculations, we use model twist-2 and twist-3 light-cone wave functions incorporating transverse degrees of freedom, where the collinear DAs are derived from QCD sum rules [34, 35, 36, 37, 38]. Naively, the twist-3 contributions to the form factor are expected to be small compared to leading (twist-2) contributions as they have a relative $1/Q^2$ suppression. On the contrary, the existing literature, either using model or asymptotic DAs [25, 27, 28, 57, 62, 64], shows large twist-3 corrections to the pion form factor which even overshoot the twist-2 contributions in a wide range of low and intermediate energies. This is also confirmed in our analysis and is in fact more enhanced for the kaon form factor. To this end, our analysis shows good agreement with the existing pion data and in addition we prove the consistency of our results by adopting a scheme of analytization of the running strong coupling [60, 61, 89, 90] that removes the explicit Landau singularity at $Q^2 = \Lambda_{\text{QCD}}^2$ by a minimum power correction in the UV regime.

The paper is organized as follows: In Section II, we briefly discuss the idea of factorization and review the basic definitions of the twist-2 and twist-3 pseudo-scalar meson DAs and their renormalization evolutions. Section III deals with the theoretical framework involved in calculating the space-like electromagnetic form factor. Here, we recall the predictions of classic asymptotic QCD for large $Q^2 \rightarrow \infty$ and how one needs to modify pQCD with collinear as well as “ k_T ” factorization schemes including Sudakov effects at intermediate energies. In Section IV, we provide the details of our numerical results for the pion form factor and compare it with the available experimental data. We also give a preliminary prediction for the kaon form factor, despite the lack of available experimental data for comparison in the desired phenomenological regime. Finally, Section V contains our summary and conclusions. The appendices contain a compendium of relevant formulae used in our analysis.

II. FACTORIZATION AND DISTRIBUTION AMPLITUDES

The parton model of describing exclusive processes in QCD inherently rest on the so-called *frozen approximation* ([70, 71, 72]). At high energies, exclusive scattering amplitudes are dominated by hadronic Fock states with essentially valence quark configurations ($\bar{q}q$ in mesons). While the relative velocities of the participating hadrons are located closely to the null-plane, the internal hadron “quantum-bindings” processes are highly time-dilated with respect to the exclusive reaction time scales in the rest frames of the remaining hadrons. This effectively freezes the hadronic internal degrees of freedom as seen by the other hadrons. This incoherence between the long-distance intra-hadronic binding processes and the short-distance inter-hadronic scattering reaction is the very motivation for the idea of factorization. Thus, the hadrons may be considered to be consisting of definite valence quark states denoted by a DA of leading twist ϕ . The collinear factorization formula is then used to express exclusive quantities like the form factors as a convolution using the DAs:

$$F_M(Q^2) = \int_0^1 dx dy \phi_{\text{in}}(x, \mu_F^2) T_H(x, y, Q^2, \mu_F^2, \mu_R^2) \phi_{\text{out}}(y, \mu_F^2) + \cdots, \quad (2)$$

where $Q^2 = -2P_{\text{in}} \cdot P_{\text{out}}$. Here, P_{in} and P_{out} are, respectively, the ingoing and outgoing hadron momenta, x and y are the longitudinal momentum fractions of the nearly on-shell valence quarks, μ_R is the renormalization scale and μ_F is the factorization scale which is defined as the scale below which the QCD dynamics are non-perturbative and remain implicitly encoded within the DAs, while the dynamics above are perturbative and must be retained in the hard kernel T_H . The ellipses in the above equation represent contributions from higher order Fock states and sub-leading twists which are all suppressed by inverse powers of Q^2 . In addition, they also include the non-factorizable soft contributions. Formally, the definition of the leading twist-2 DA for pseudo-scalar mesons (e.g., π^-) can be given in a process- and frame-independent manner [34, 35, 70, 71, 72] in terms of matrix elements of non-local light-ray operator along a certain light-like direction z_μ ($z^2 = 0$):

$$\langle 0 | \bar{u}(z)[z, -z]\gamma_\mu\gamma_5 d(-z) | \pi^-(P) \rangle = iP_\mu \int_0^1 dx e^{i\xi(zP)} \phi_{2,\pi}(x, \mu_F^2); \xi = 2x - 1, \quad (3)$$

with the path-ordering ($\mathcal{P}$) Wilson line in terms of the gluon “connection” along the straight line joining z and $-z$ along the null-plane given by

$$[z, -z] = \mathcal{P} \left[ig_s \int_{-z}^z dy^\mu A_\mu(y) \right], \quad (4)$$

where $P_\mu^2 = m_\pi^2$ and p_μ is a light-like vector,

$$p_\mu = P_\mu - \frac{1}{2} z_\mu \frac{m_\pi^2}{Pz}. \quad (5)$$

The local limit $z \rightarrow 0$ gives the normalization condition at an arbitrary scale μ ,

$$\int_0^1 \phi_{2;\pi}(x, \mu^2) dx = \frac{f_\pi}{2\sqrt{2N_c}} \quad (6)$$

with the pion decay constant, $f_\pi \approx 131$ MeV defined by

$$\langle 0 | \bar{u}(0) \gamma_\mu \gamma_5 d(0) | \pi^-(P) \rangle = i f_\pi P_\mu. \quad (7)$$

The leading twist-2 DA $\phi_{2;\pi}(x, \mu^2)$ can be expressed as a conformal series expansion over Gegenbauer polynomials $C_{2n}^{3/2}$:

$$\phi_{2;\pi}(x, \mu^2) = \frac{3f_\pi}{\sqrt{2N_c}} x(1-x) \left(1 + \sum_{n=1}^{\infty} a_{2n}^\pi(\mu^2) C_{2n}^{3/2}(\xi) \right), \quad (8)$$

where

$$\phi_{2;\pi}^{(\text{as})}(x) = \phi_{2;\pi}(x, \mu^2 \rightarrow \infty) = \frac{3f_\pi}{\sqrt{2N_c}} x(1-x) \quad (9)$$

is generally referred to as the *asymptotic* DA. The Gegenbauer moments a_{2n}^π represent the non-perturbative inputs encoding the long-distance dynamics and may be obtained e.g., via lattice QCD calculations or QCD sum rules. The renormalization group (RG) equation for $\phi_{2;\pi}(x, \mu^2)$ is known as the Efremov-Radyushkin-Brodsky-Lepage (ER-BL) equation [70, 71, 72],

$$\mu^2 \frac{d}{d\mu^2} \phi_{2;\pi}(x, \mu^2) = \int_0^1 dy V(x, y; \alpha_s(\mu^2)) \phi_{2;\pi}(y, \mu^2) \quad (10)$$

with the integral kernel $V(x, y; \alpha_s)$ to leading order in α_s given by

$$V_0(x, y; \alpha_s) = \mathcal{C}_F \frac{\alpha_s}{2\pi} \left[\frac{1-x}{1-y} \left(1 + \frac{1}{x-y} \right) \theta(x-y) + \frac{x}{y} \left(1 + \frac{1}{y-x} \right) \theta(y-x) \right]_+, \quad (11)$$

where the “+” distribution is defined as

$$[V(x, y; \alpha_s)]_+ = V(x, y; \alpha_s) - \delta(x-y) \int_0^1 dt V(t, y; \alpha_s). \quad (12)$$

Solving the above set of equations yields the multiplicative renormalization formula for moments a_n^π to leading-logarithmic accuracy,

$$a_n(\mu^2) = L^{\gamma_n^{(0)}/\beta_0} a_n(\mu_0^2), \quad (13)$$

where $L = \alpha_s(\mu^2)/\alpha_s(\mu_0^2)$ and $\beta_0 = (11N_c - 2N_f)/12$, while the lowest order anomalous dimensions are given by

$$\gamma_n^{(0)} = \mathcal{C}_F \left(\psi(n+2) + \psi(1) - \frac{3}{4} - \frac{1}{2(n+1)(n+2)} \right) \quad (14)$$

with the logarithmic derivative of the Gamma function $\psi(z) = \Gamma'(z)/\Gamma(z)$. Note that for the pion all odd moments $a_{n=1,3,5,\dots}^\pi$ vanish due to isospin symmetry. In contrast, the kaon DA have non-zero values for the odd moments signifying flavor-SU(3) violation effects. Hence, the twist-2 DA for the kaon (e.g., K^-) is given by the expansion

$$\phi_{2;K}(x, \mu^2) = \frac{3f_K}{\sqrt{2N_c}} x(1-x) \left(1 + \sum_{n=1}^{\infty} a_n^K(\mu^2) C_n^{3/2}(\xi) \right); \quad (15)$$

$$\int_0^1 \phi_{2;K}(x, \mu^2) dx = \frac{f_K}{2\sqrt{2N_c}}, \quad (16)$$

where the kaon decay constant $f_K \approx 1.22f_\pi$ [24] is defined by

$$\langle 0 | \bar{u}(0) \gamma_\mu \gamma_5 s(0) | K^-(P) \rangle = i f_K P_\mu. \quad (17)$$

The Gegenbauer moments being multiplicatively renormalizable with growing anomalous dimensions, for sufficiently large renormalization scale a finite number of moments are relevant, albeit the fact that the higher order moments have large uncertainties in their present determination. Hence, in all practical calculations, the series expansion of the DAs are truncated only to the first few moments. In this paper, we have adopted a model for the twist-2 DAs in truncating up to the second moment, as was done in Refs. [35, 37].

For the charged pseudo-scalar mesons at the twist-3 level, there are two 2-particle DAs and one 3-particle DA. Here, we only give the formal definitions of the 2-particle DAs that we need in our analysis. For the charged pion (e.g., π^-), they are defined by [34] :

$$\begin{aligned} \langle 0 | \bar{u}(z) i \gamma_5 d(-z) | \pi^-(P) \rangle &= \mu_\pi \int_0^1 dx e^{i\xi(zp)} \phi_{3;\pi}^p(x, \mu^2), \\ \langle 0 | \bar{u}(z) \sigma_{\alpha\beta} \gamma_5 d(-z) | \pi^-(P) \rangle &= -\frac{i}{3} \mu_\pi (P_\alpha z_\beta - P_\beta z_\alpha) \int_0^1 dx e^{i\xi(zp)} \phi_{3;\pi}^\sigma(x, \mu^2) \end{aligned} \quad (18)$$

with $\mu_\pi = m_\pi^2/(m_u + m_d)$ and similarly for the charged kaon (e.g., K^-) [37] :

$$\begin{aligned} \langle 0 | \bar{u}(z) i \gamma_5 s(-z) | K^-(P) \rangle &= \mu_K \int_0^1 dx e^{i\xi(zp)} \phi_{3;K}^p(x, \mu^2), \\ \langle 0 | \bar{u}(z) \sigma_{\alpha\beta} \gamma_5 s(-z) | K^-(P) \rangle &= -\frac{i}{3} \mu_K (P_\alpha z_\beta - P_\beta z_\alpha) \int_0^1 dx e^{i\xi(zp)} \phi_{3;K}^\sigma(x, \mu^2) \end{aligned} \quad (19)$$

with $\mu_K = m_K^2/(m_u + m_s)$. Note that the gauge-link factors (Wilson line (4)) in the matrix elements are to be implicitly understood. The twist-3 DAs have the following asymptotic forms:

$$\begin{aligned}\phi_{3;M}^{p(\text{as})}(x) &= \frac{f_M}{4\sqrt{2N_c}}, \\ \phi_{3;M}^{\sigma(\text{as})}(x) &= \frac{3f_M}{2\sqrt{2N_c}} x(1-x) ; \quad M = \pi^\pm, K^\pm\end{aligned}\tag{20}$$

with the normalization condition,

$$\int_0^1 \phi_{3;M}^{p,\sigma}(x, \mu^2) = \frac{f_M}{4\sqrt{2N_c}}.\tag{21}$$

For our analysis, we use the 2-particle twist-3 DAs from Refs. [35, 37] defined at the scale $\mu = 1$ GeV. As a matter of book-keeping, we explicitly provide the relevant formulae for the charged pion and kaon DAs in Appendix A.

III. SPACE-LIKE ELECTROMAGNETIC FORM FACTOR

The electromagnetic form-factor is considered as an important observable for studying the onset of the perturbative regime in exclusive processes. For large Q^2 , the asymptotic scaling behavior $F_M(Q^2) \sim 1/Q^2$ follows from the well-known dimensional “quark counting” while for small Q^2 , the behavior is well described by the Vector Meson Dominance (VMD) model [42, 43, 44] and given by

$$F_M(Q^2) \approx \frac{1}{1 + Q^2/\mu_{\text{VDM}}^2} ; \quad Q^2 \ll \mu_{\text{VDM}}^2,\tag{22}$$

where $\mu_{\text{VDM}} \approx 750$ MeV is a reasonable cut-off mass scale, showing no obvious trace of pQCD scaling behavior where there exists no high energy cut-off. Hence, a thorough understanding of this transition behavior (from nonperturbative to perturbative) is of crucial importance in QCD for understanding the very nature of strong interaction and in providing a vivid picture of the underlying quark-gluon substructure of the mesons.

For a charged meson M (e.g., $\pi^\pm, K^\pm$), the form factor is specified by the following matrix element:

$$(P' + P)_\mu F_M(Q^2) = \langle M(P') | J_\mu(0) | M(P) \rangle ; \quad J_\mu = \sum_f e_f \bar{q}_f \gamma_\mu q_f,\tag{23}$$

where J_μ is the electromagnetic current with quark q_f of flavor f and charge e_f . In this paper, we shall only consider space-like momentum transfers i.e., $q^2 = (P' - P)^2 = -Q^2$.

Neglecting the meson masses, we consider the “brick wall” frame where the incoming particle with 4-momentum P in the z direction recoils with 4-momentum P' in the $-z$ direction after interacting with the hard photon “wall”. In the light-cone formalism, $P = (Q/\sqrt{2}, 0, \mathbf{0}_T)$ and $P' = (0, Q/\sqrt{2}, \mathbf{0}_T)$.

A. Hard contribution in pQCD

The hard contributions to the form factor are calculated using the collinear factorization formula Eq. (2), where the hard scattering kernel T_H at the scale $\mu = \mu_F = \mu_R$ is given to the leading order in α_s by

$$T_H(x, y, Q^2, \mu^2) = 16\pi\mathcal{C}_F\alpha_s(\mu^2) \left[\frac{2}{3} \frac{1}{xyQ^2} + \frac{1}{3} \frac{1}{(1-x)(1-y)Q^2} \right], \quad (24)$$

where in QCD the value of the *Casimir* operator in the fundamental representation of SU(3) is $\mathcal{C}_F = (N_c^2 - 1)/2N_c = 4/3$. The factorization formula then yields the classic pQCD expression for the meson form factor at $\mu^2 = Q^2$:

$$F_M^{\text{hard}}(Q^2) = \frac{16\pi\mathcal{C}_F\alpha_s(Q^2)}{Q^2} \left| \int_0^1 dx \frac{\phi_{2;M}(x, Q^2)}{x} \right|^2. \quad (25)$$

Note that using the asymptotic twist-2 DA $\phi_{2;M}^{(\text{as})}(x)$, one obtains the familiar $1/Q^2$ scaling behavior for $Q^2 \rightarrow \infty$

$$F_M^{\text{hard}}(Q^2) = \frac{8\pi\alpha_s(Q^2)f_M^2}{Q^2}. \quad (26)$$

The principal motivation of the modified “resummed” pQCD is the elimination of large logarithms in the hard kernel that arise from radiative gluon loop corrections. One way of doing this is by the introduction of intrinsic transverse momenta dependence of the constituent partons, giving rise to a Sudakov suppression due to certain partial resummation of transverse terms, as mentioned earlier in the introduction. Including the transverse momenta of the two valence quarks within the meson, the tree-level hard kernel T_H in the momentum-space is written as

$$T_H(x, y, Q^2, \mathbf{k}_{1T}, \mathbf{k}_{2T}, \mu^2) = \frac{16\pi\mathcal{C}_F\alpha_s(\mu^2)xQ^2}{(xQ^2 + \mathbf{k}_{1T}^2)(xyQ^2 + (\mathbf{k}_{1T} - \mathbf{k}_{2T})^2)}, \quad (27)$$

where the transverse momentum dependence now sets the factorization scale. Then the modified factorization formula in the transverse *impact parameter* representation is given by

$$F_M^{\text{hard}}(Q^2) = \int_0^1 dx dy \int \frac{d^2b_1}{(2\pi)^2} \frac{d^2b_2}{(2\pi)^2} \times \mathcal{P}_{2;M}(x, b_1, P, \mu) \tilde{T}_H(x, y, Q, b_1, b_2, \mu) \mathcal{P}_{2;M}(y, b_2, P', \mu), \quad (28)$$

where the modified DA $\mathcal{P}_{2;M}(x_i, b_i, P_i, \mu)$ absorbs the large infrared logarithms into the Sudakov exponent S_i [22] (including also the evolution of the DA from the factorization scale $1/b_i$ to the scale μ):

$$\begin{aligned}\mathcal{P}_{2;M}(x_i, b_i, P_i \simeq Q, \mu) &= \exp[-S_i(XQ, b_i, \mu)] \tilde{\mathcal{P}}_{2;M}(x_i, b_i, 1/b_i); \\ S_i(XQ, b_i, \mu) &= s(x_i Q, 1/b_i) + s((1-x_i)Q, 1/b_i) + 2 \int_{1/b_i}^{\mu} \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu}^2)), \\ s(XQ, 1/b_i) &= \int_{1/b_i}^{XQ/\sqrt{2}} \frac{d\mu}{\mu} \left[\ln\left(\frac{XQ}{\sqrt{2}\mu}\right) \mathcal{A}(\alpha_s(\mu^2)) + \mathcal{B}(\alpha_s(\mu^2)) \right],\end{aligned}\quad (29)$$

where $1/b_1, 1/b_2$ set the factorization scales in the transverse impact configuration. In the above equations, the quark anomalous dimension is given by $\gamma_q(\alpha_s) = -\alpha_s/\pi$ and the “cusp” anomalous dimensions $\mathcal{A}$ and $\mathcal{B}$, to one-loop accuracy are given by

$$\begin{aligned}\mathcal{A}(\alpha_s(\mu^2)) &= \mathcal{C}_F \frac{\alpha_s(\mu^2)}{\pi} + \left[\left(\frac{67}{27} - \frac{\pi^2}{9} \right) N_c - \frac{10}{27} N_f + \frac{8}{3} \beta_0 \ln\left(\frac{e^{\gamma_E}}{2}\right) \right] \left(\frac{\alpha_s(\mu^2)}{\pi} \right)^2, \\ \mathcal{B}(\alpha_s(\mu^2)) &= \frac{2}{3} \frac{\alpha_s(\mu^2)}{\pi} \ln\left(\frac{e^{2\gamma_E-1}}{2}\right)\end{aligned}\quad (30)$$

where the $\overline{\text{MS}}$ running coupling to two-loop accuracy in standard perturbation theory is given by

$$\frac{\alpha_s(\mu^2)}{\pi} = \frac{1}{\beta_0 \ln(\mu^2/\Lambda_{\text{QCD}}^2)} - \frac{\beta_1 \ln(\ln(\mu^2/\Lambda_{\text{QCD}}^2))}{\beta_0^3 \ln^2(\mu^2/\Lambda_{\text{QCD}}^2)} \quad (31)$$

with $\beta_0 = (11N_c - 2N_f)/12 = 9/4$ and $\beta_1 = (51N_c - 19N_f)/24 = 4$ for $N_c = N_f = 3$. Note that the above modified factorization calls for introducing a scale hierarchy $XQ > 1/b_i > \Lambda_{\text{QCD}}$ (where $X = x_i, (1-x_i)$, $x_1 = x$ and $x_2 = y$) to separate the distinct contributions from the perturbative and non-perturbative kinematic regions without the possibility of a “double counting”. Note that there exist other schemes of defining the running coupling involving power corrections, restoring the explicit Landau singularity and the analyticity at $Q^2 = 0$ (see e.g., Refs. [89, 90] and also Section IV for details.)

At low momentum transfers, the modified infrared free DAs are often approximated with constituent quark masses which are different from the actual masses of the current quarks and usually chosen close to the intrinsic transverse scale Λ_{QCD} of the hadron structure, i.e., between 200-500 MeV. These quark masses which effectively parametrize the QCD vacuum effects are also used to suppress possible endpoint effects. Hence, we have

$$\tilde{\mathcal{P}}_{2;M}(x_i, b_i, 1/b_i) \simeq \tilde{\mathcal{P}}_{2;M}(x_i, b_i, 1/b_i, \mathcal{M}_q) \quad (32)$$

which could be expressed in terms of the full momentum-space light-cone wave function $\Psi_{2;M}$ (which also includes the transverse momentum distribution of the constituent bound state partons):

$$\tilde{\mathcal{P}}_{2;M}(x_i, b_i, 1/b_i, \mathcal{M}_q) = \int_{\mathbf{k}_{iT}^2 \leq (1/b_i)^2} \frac{d^2 \mathbf{k}_{iT}}{16\pi^3} \Psi_{2;M}(x_i, \mathbf{k}_{iT}, 1/b_i, \mathcal{M}_q). \quad (33)$$

To model the intrinsic transverse momentum dependence of the meson wave functions, we use the Brodsky-Huang-Lepage (BHL) gaussian prescription [81, 87]:

$$\Psi_{2;M}(x_i, \mathbf{k}_{iT}, 1/b_i, \mathcal{M}_q) = \Phi_{2;M}(x_i, 1/b_i) \Sigma(x_i, \mathbf{k}_{iT}, \mathcal{M}_q) \quad (34)$$

with

$$\Phi_{2;M}(x_i, 1/b_i) = A_{2;M} \phi_{2;M}(x_i, 1/b_i), \quad (35)$$

$$\Sigma(x_i, \mathbf{k}_{iT}, \mathcal{M}_q) = \frac{16\pi^2 \beta_{2;M}^2}{x_i(1-x_i)} \exp \left[-\frac{\beta_{2;M}^2}{x_i(1-x_i)} (\mathbf{k}_{iT}^2 + \mathcal{M}_q^2) \right], \quad (36)$$

assuming equal masses of the two constituent quarks within the meson. The parameters $A_{2;M}$, $\beta_{2;M}$ and $\mathcal{M}_q$ are fixed using phenomenological constraints. The above integration then yields the full modified wave function in the impact representation:

$$\begin{aligned} \tilde{\mathcal{P}}_{2;M}(x_i, b_i, 1/b_i, \mathcal{M}_q) &= A_{2;M} \phi_{2;M}(x_i, 1/b_i) \exp \left[-\frac{\beta_{2;M}^2 \mathcal{M}_q^2}{x_i(1-x_i)} \right] \\ &\quad \times \exp \left[-\frac{b_i^2 x_i(1-x_i)}{4\beta_{2;M}^2} \right] \end{aligned} \quad (37)$$

Including the RG evolution equation for the hard kernel:

$$\tilde{T}_H(x, y, Q, b_1, b_2, \mu) = \exp \left[-4 \int_\mu^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu}^2)) \right] \tilde{T}_H(x, y, Q, b_1, b_2, t), \quad (38)$$

where

$$t = \max(\sqrt{xy} Q, 1/b_1, 1/b_2), \quad (39)$$

one arrives at the “double-b” factorization formula for the meson form factor at the twist-2 level [58] :

$$\begin{aligned} F_M^{(t=2)}(Q^2) &= 16\pi Q^2 \mathcal{C}_F \int_0^1 x dx dy \int_0^\infty b_1 db_1 b_2 db_2 \alpha_s(t) \tilde{\mathcal{P}}_{2;M}(x, b_1, 1/b_1, \mathcal{M}_q) \\ &\quad \times \tilde{\mathcal{P}}_{2;M}(y, b_2, 1/b_2, \mathcal{M}_q) H(x, y, Q, b_1, b_2) \exp[-S(x, y, b_1, b_2, Q)] \end{aligned} \quad (40)$$

with

$$H(x, y, Q, b_1, b_2) = K_0(\sqrt{xy} Q b_2) \left[\theta(b_1 - b_2) K_0(\sqrt{x} Q b_1) I_0(\sqrt{x} Q b_2) \right. \\ \left. + \theta(b_2 - b_1) K_0(\sqrt{x} Q b_2) I_0(\sqrt{x} Q b_1) \right]. \quad (41)$$

K_0 and I_0 are modified Bessel functions and the full Sudakov exponent is given by

$$S(x, y, b_1, b_2, Q) = \sum_{i=1}^2 \left[s(x_i Q, 1/b_i) + s((1 - x_i) Q, 1/b_i) + 2 \int_{1/b_i}^t \frac{d\bar{\mu}}{\bar{\mu}} \gamma_q(\alpha_s(\bar{\mu}^2)) \right]. \quad (42)$$

For completeness, the expression for the full Sudakov factor $s(XQ, 1/b_i)$, up to next-to-leading logarithm accuracy is given in the Appendix B. The expression slightly differs from the result given in Ref. [91], but numerically this difference is insignificant at our working accuracy. Note that this difference was first observed in Ref. [60].

To include the sub-leading twist-3 corrections to the form factor, the hard scattering kernel gets slightly modified as compared to the twist-2 case which turns out to be [57]

$$T_H^{(t=3)}(x, y, Q^2, \mathbf{k}_{1T}, \mathbf{k}_{2T}, \mu^2) = \frac{64\pi \mathcal{C}_F \alpha_s(\mu^2) x}{(xQ^2 + \mathbf{k}_{1T}^2)(xy Q^2 + (\mathbf{k}_{1T} - \mathbf{k}_{2T})^2)}. \quad (43)$$

Applying the momentum projection operator [92, 93],

$$\mathcal{M}_{\alpha\beta}^M = i \left\{ \not{P} \gamma_5 \Psi_{2,M} - \mu_M \gamma_5 \left(\Psi_{3;M}^p - i \sigma_{\mu\nu} n^\mu \bar{n}^\nu \frac{\Psi_{3;M}^{\sigma'}}{6} + i \sigma_{\mu\nu} P^\mu \frac{\Psi_{3;M}^\sigma}{6} \frac{\partial}{\partial k_{T\nu}} \right) \right\}_{\alpha\beta} \quad (44)$$

on the bilocal matrix element with quark flavors f_1 and f_2 ($f_{1,2} = u, d, s$),

$$\langle 0 | \bar{q}_{f_1}(z) q_{f_2}(-z) | M(P) \rangle_{\alpha\beta} = i \int_0^1 dx \int d^2 \mathbf{k}_T e^{i\xi(zp)} \\ \times \left\{ \not{P} \gamma_5 \Psi_{2;M} - \mu_M \gamma_5 \left(\Psi_{3;M}^p - \sigma_{\mu\nu} P^\mu z^\nu \frac{\Psi_{3;M}^\sigma}{6} \right) \right\}_{\alpha\beta}, \quad (45)$$

where $\Psi_{3;M}^{\sigma'}(x, \mathbf{k}_T, 1/b, \mathcal{M}_q) = \partial \Psi_{3;M}^\sigma(x, \mathbf{k}_T, 1/b, \mathcal{M}_q) / \partial x$, $n = (1, 0, \mathbf{0}_T)$ is the unit vector in the “+” direction, $\bar{n} = (0, 1, \mathbf{0}_T)$ is the unit vector in the “-” direction, $\xi = 2x - 1$ and $\mu_M = m_M^2 / (m_{q_{f_1}} + m_{q_{f_2}})$, one obtains the final formula for hard meson form factor up to

twist-3 corrections given by [62, 64]

$$\begin{aligned}
F_M^{\text{hard}}(Q^2) &= F_M^{(t=2)}(Q^2) + F_M^{(t=3)}(Q^2) \\
&= 32\pi Q^2 \mathcal{C}_F \int_0^1 dx dy \int_0^\infty b_1 db_1 b_2 db_2 \alpha_s(t) \\
&\quad \times \left[\frac{x}{2} \tilde{\mathcal{P}}_{2;M}(x, b_1, 1/b_1, \mathcal{M}_q) \tilde{\mathcal{P}}_{2;M}(y, b_2, 1/b_2, \mathcal{M}_q) \right. \\
&\quad + \frac{\mu_M^2}{Q^2} \left(\bar{x} \tilde{\mathcal{P}}_{3;M}^p(x, b_1, 1/b_1, \mathcal{M}_q) \tilde{\mathcal{P}}_{3;M}^p(y, b_2, 1/b_2, \mathcal{M}_q) \right. \\
&\quad + \frac{(1+x)}{6} \tilde{\mathcal{P}}_{3;M}^p(x, b_1, 1/b_1, \mathcal{M}_q) \tilde{\mathcal{P}}_{3;M}^{\sigma'}(y, b_2, 1/b_2, \mathcal{M}_q) \\
&\quad \left. \left. + \frac{1}{2} \tilde{\mathcal{P}}_{3;M}^p(x, b_1, 1/b_1, \mathcal{M}_q) \tilde{\mathcal{P}}_{3;M}^\sigma(y, b_2, 1/b_2, \mathcal{M}_q) \right) \right] H(x, y, Q, b_1, b_2) \\
&\quad \times \prod_{i=1}^2 S_t(x_i) S_t(\bar{x}_i) \exp[-S(x, y, b_1, b_2, Q)] \ ; \ \bar{x}_i = 1 - x_i \ .
\end{aligned} \tag{46}$$

Here, we have assumed a similar gaussian ansatz in the transverse momentum distribution of the modified twist-3 wave functions:

$$\begin{aligned}
\tilde{\mathcal{P}}_{3;M}^p(x_i, b_i, 1/b_i, \mathcal{M}_q) &= A_{3;M}^p \phi_{3;M}^p(x_i, 1/b_i) \exp \left[-\frac{(\beta_{3;M}^p)^2 \mathcal{M}_q^2}{x_i(1-x_i)} \right] \\
&\quad \times \exp \left[-\frac{b_i^2 x_i(1-x_i)}{4(\beta_{3;M}^p)^2} \right] \ , \\
\tilde{\mathcal{P}}_{3;M}^\sigma(x_i, b_i, 1/b_i, \mathcal{M}_q) &= A_{3;M}^\sigma \phi_{3;M}^\sigma(x_i, 1/b_i) \exp \left[-\frac{(\beta_{3;M}^\sigma)^2 \mathcal{M}_q^2}{x_i(1-x_i)} \right] \\
&\quad \times \exp \left[-\frac{b_i^2 x_i(1-x_i)}{4(\beta_{3;M}^\sigma)^2} \right] \ .
\end{aligned} \tag{47}$$

The hard kernel H and the Sudakov exponent S are given by Eq. (41) and Eq. (42), respectively. The above formula is used to evaluate the pion and kaon hard form factors using the twist-2 DAs Eq. (8) and Eq. (15), respectively, and twist-3 DAs provided in Appendix A. The $S_t(x_i)$ are *jet functions*, defined as eikonalized matrix elements of quark fields attached by a Wilson line, arising from another kinematic resummation scheme called the *threshold resummation*, as introduced in Ref. [94, 95]. The modified treatment of the collinear factorization prescription works reasonably well for the twist-2 case but for the twist-3 case, the Sudakov suppression factor may still not be effective enough in shielding the non-perturbative enhancements due to endpoint singularities. These are kinematic singularities of the scattering amplitude when the longitudinal momentum fraction x of the

valence partons (quarks) go to 0, 1. Therefore, in addition there is a need to sum up the collinear double logarithms of the type $\alpha_s \ln^2 x$ to all orders, which are then collected into these jet functions. The exact form of $S_t(x_i)$ involves a one parameter integration, but for the sake of numerical calculations it is convenient to take the simple parameterization, as proposed in Ref. [94, 95] :

$$S_t(x_i) = \frac{2^{1+2c}\Gamma(3/2+c)}{\sqrt{\pi}\Gamma(1+c)} [x_i(1-x_i)]^c, \quad (48)$$

where the parameter $c \approx 0.3$ for light pseudo-scalar mesons like the pion and kaon. The jet functions vanish at the endpoints and modify the endpoint behavior of the DAs, providing enough suppression to damp the artificial effect of endpoint singularities.

B. Soft contributions via Local Duality

The perturbative predictions for the pion form factor are known to be relatively small for phenomenological low momentum transfers ($Q^2 \leq 10 \text{ GeV}^2$) [48, 49, 50, 69, 78, 83, 85], as also evident from our analysis in the next section. Clearly, there is the need for including non-factorizable soft contributions to explain the experimental data. The factorization ansatz Eq. (2) holds for large momentum transfers under the assumption that only the contributions from valence parton states dominate. This approximation no longer holds true at small momenta when contributions from higher Fock states with more than valence partons become significant. In addition, there could be non-perturbative enhancements from the so-called *Feynman mechanism*, which corresponds to selecting a hadronic configuration in which one of the valence parton carries almost the entire hadron momenta. Unfortunately, due to the complexity of soft QCD processes, there are no unambiguous ways to calculate these contributions analytically using the parton picture and Feynman diagrammatics, other than using theoretical models for the DAs. In this paper, we follow the Local Duality (LD) approach from QCD sum rules as in Ref. [61] where the same problem is addressed without a direct reference to DAs. In this section, we simply use the result for the soft form factor derived in the LD approach:

$$F_M^{\text{soft}}(Q^2) = F_M^{\text{LD}}(Q^2) = 1 - \frac{1 + 6s_0(Q^2)/Q^2}{(1 + 4s_0(Q^2)/Q^2)^{3/2}}. \quad (49)$$

The *duality interval* s_0 , encodes the non-perturbative information about higher excited states and continuum contributions and is given by

$$s_0(Q^2) = 4\pi^2 f_M^2 / (1 + \frac{\alpha_s(Q^2)}{\pi}). \quad (50)$$

Expanding in inverse powers of Q gives $F_M^{\text{soft}}(Q^2) \sim 1/Q^4$ for large Q^2 and is thus expected to be sub-leading compared to the leading perturbative contribution from Eq. (26). Nevertheless, at low and moderate momentum transfers the soft contributions turn out to be so significant in obtaining a good agreement with the experimental data. This fact is clearly revealed in our analysis in the next section.

Next we add together the hard and the soft contribution to obtain the total contributions to the electromagnetic form factor $F_M(Q^2)$. Here, it is necessary to ensure that the respective contributions lie within their domains of validity to minimize the possibility of a double counting. This technique, as introduced in Ref. [61], employs gauge invariance that protects the value $F_M(0) = 1$, through the vector Ward identity relating a 3-point Green function to a 2-point Green function at zero momentum transfer, i.e., $F_M^{\text{LD}}(Q^2=0) = 1$. This implies that $F_M^{\text{hard}}(Q^2=0) = 0$. A “smooth” transition from the hard to the soft behavior is then ensured by a *matching ansatz* from the large Q^2 behavior (arising from $F_M^{\text{hard}}(Q^2)$) to the low Q^2 behavior (arising from $F_M^{\text{soft}}(Q^2)$). This can be done by introducing a mass scale M_0 which in the LD approach should be identified with the threshold $M_0^2 = 2s_0$. The twist-2 part of the hard form factor $F_M^{(t=2)}(Q^2)$ is then modified following Ref. [61]

$$F_M^{(t=2)}(Q^2) \rightarrow \left(\frac{Q^2}{2s_0(Q^2) + Q^2} \right)^2 F_M^{(t=2)}(Q^2). \quad (51)$$

However, for the twist-3 case, the “matching function” $\Phi(z) = 1/(1+z)^2$, with $z = Q^2/M_0^2$ is insufficient to ensure the Ward identity at $Q^2 = 0$. To correct for the singular ($\sim 1/Q^4$) behavior, we make a similar modification of the twist-3 part via the replacement

$$F_M^{(t=3)}(Q^2) = \tilde{F}_M^{(t=3)}(Q^2) \frac{M_0^4}{Q^4} \rightarrow \tilde{F}_M^{(t=3)}(Q^2) \frac{M_0^4}{M_0^4 + Q^4} \quad (52)$$

with the choice of the matching function $\tilde{\Phi}(z) = 1/(1+z^2)^2$. This yields the Ward identity corrected twist-3 part:

$$F_M^{(t=3)}(Q^2) \rightarrow \left(\frac{Q^4}{4s_0^2(Q^2) + Q^4} \right)^2 F_M^{(t=3)}(Q^2). \quad (53)$$

Finally, we arrive at our expression for the total electromagnetic form factor for a charged meson M ($\pi^\pm, K^\pm$), valid for all values of Q^2 and is given by

$$F_M(Q^2) = 1 - \frac{1 + 6s_0(Q^2)/Q^2}{(1 + 4s_0(Q^2)/Q^2)^{3/2}} + \left(\frac{Q^2}{2s_0(Q^2) + Q^2} \right)^2 F_M^{(t=2)}(Q^2) + \left(\frac{Q^4}{4s_0^2(Q^2) + Q^4} \right)^2 F_M^{(t=3)}(Q^2), \quad (54)$$

where $F_M^{(t=2)}(Q^2)$ and $F_M^{(t=3)}(Q^2)$ are given from Eq. (40) and Eq. (46), respectively.

IV. NUMERICAL RESULTS

At first, we need to determine the pion and kaon gaussian parameters $A_{2;M}$, $A_{3;M}^p$, $A_{3;M}^\sigma$ and $\beta_{2;M}$, $\beta_{3;M}^p$, $\beta_{3;M}^\sigma$ for the twist-2 and twist-3 light-cone wave functions, respectively. For the pion, they are obtained from the two constraints: firstly, by virtue of the leptonic decay $\pi \rightarrow \mu\nu_\mu$, we have the condition

$$\int_0^1 dx \int \frac{d^2\mathbf{k}_T}{16\pi^3} \Psi_\pi(x, \mathbf{k}_T, \mathcal{M}_{u,d}) = \frac{f_\pi}{2\sqrt{2N_c}}, \quad (55)$$

leading to

$$A_\pi \int_0^1 dx \phi_\pi(x) \exp \left[-\frac{\beta_\pi^2 \mathcal{M}_{u,d}^2}{x(1-x)} \right] = \frac{f_\pi}{2\sqrt{6}} \quad (56)$$

and secondly, from $\pi^0 \rightarrow \gamma\gamma$, we have the condition

$$\int_0^1 dx \Psi_\pi(x, \mathbf{k}_T=0, \mathcal{M}_{u,d}) = \frac{\sqrt{2N_c}}{f_\pi}, \quad (57)$$

which implies

$$16A_\pi\beta_\pi^2\pi^2 \int_0^1 dx \frac{\phi_\pi(x)}{x(1-x)} \exp \left[-\frac{\beta_\pi^2 \mathcal{M}_{u,d}^2}{x(1-x)} \right] = \frac{\sqrt{6}}{f_\pi}, \quad (58)$$

where we use the constituent quark mass $\mathcal{M}_{u,d} = 0.33$ GeV for both the u and d valence quarks in the pion. In the case of the kaon, firstly from the leptonic decay $K \rightarrow \mu\nu_\mu$, we have the constraint

$$A_K \int_0^1 dx \phi_K(x) \exp \left[-\beta_K^2 \left(\frac{\mathcal{M}_s^2}{x} + \frac{\mathcal{M}_{u,d}^2}{1-x} \right) \right] = \frac{f_K}{2\sqrt{6}}. \quad (59)$$

As for the second constraint, no straightforward condition like Eq. (57) could be obtained for the kaon. On the other hand, by virtue of SU(3) isospin symmetry, it is reasonable to

make an assumption that for the kaon the average transverse momentum-squared of the valence partons defined by

$$\langle \mathbf{k}_T^2 \rangle_K = \frac{\int dx \int d^2 \mathbf{k}_T |\mathbf{k}_T^2| |\Psi_K(x, \mathbf{k}_T, \mathcal{M}_{u,d,s})|^2}{\int dx \int d^2 \mathbf{k}_T |\Psi_K(x, \mathbf{k}_T, \mathcal{M}_{u,d,s})|^2}, \quad (60)$$

has about the same value as in the case of the pion. We have checked that for both the twist-2 and twist-3 pion wave functions $\langle \mathbf{k}_T^2 \rangle_\pi^{1/2} \approx 0.35$ GeV. This yields our second condition for determining the wave function parameters:

$$(0.35)^2 \approx \frac{1}{2\beta_K^2} \frac{\int_0^1 dx \phi_K^2(x) \exp \left[-2\beta_K^2 \left(\frac{\mathcal{M}_s^2}{x} + \frac{\mathcal{M}_{u,d}^2}{1-x} \right) \right]}{\int_0^1 dx \frac{\phi_K^2(x)}{x(1-x)} \exp \left[-2\beta_K^2 \left(\frac{\mathcal{M}_s^2}{x} + \frac{\mathcal{M}_{u,d}^2}{1-x} \right) \right]}, \quad (61)$$

where $\mathcal{M}_s = 0.45$ GeV is used as the constituent s -quark mass and the full light-cone kaon wave function is given by

$$\Psi_K(x, \mathbf{k}_T, \mathcal{M}_{u,d,s}) = \frac{16\pi^2 \beta_K^2 A_K}{x(1-x)} \phi_K(x) \exp \left[-\beta_K^2 \left(\frac{\mathbf{k}_T^2 + \mathcal{M}_s^2}{x} + \frac{\mathbf{k}_T^2 + \mathcal{M}_{u,d}^2}{1-x} \right) \right] \quad (62)$$

with x being the longitudinal momentum fraction of the s quark. For our numerical analysis we use typical “double-humped” type [70, 71, 72] DAs $\phi_{\pi,K}(x, \mu^2)$, derived in the framework of QCD sum rules [34, 35, 37]. Note that we have considered $N_c = 3$ in the expressions for the DAs. In Fig. 1 and Fig. 2, we display the twist-2 and twist-3 light-cone wave functions for the pion and kaon, respectively, along with their corresponding asymptotic wave functions. Note that the plots exclude the normalization factors of $\frac{1}{2\sqrt{6}}$ and $\frac{1}{4\sqrt{6}}$ for the individual DAs to facilitate comparison with one another. All the DAs are defined at the scale $\mu_0 = 1$ GeV. The twist-2 and twist-3 DA input parameters are taken from Table 3 of Ref. [37] which we again provide in Table I along with the rest of the input parameters for the wave functions. Note that for the kaon we have shown both the type of wave functions, i.e., with and without including the G-parity-breaking terms.

In Refs. [34, 35, 37], the DAs were assumed to obey equations of motion (EOM) of on-shell quarks for which $\mu_M = m_M^2 / (m_q + m_{q,s}) \approx 1.7$ GeV was used. This is not strictly correct, since the quarks are not exactly on-shell but instead confined within the hadrons. We therefore prefer using a “chiral-enhancement” parameter $\chi_{3M}(1\text{GeV}) \approx \mu_\pi \approx \mu_K$, instead of μ_M in both the DAs and also the expression for the hard form factor Eq. (46). Its numerical value is fixed by fitting the total form factor Eq. (54) to the available “world-data” for the pion [1, 2, 3, 4, 5, 6, 7, 8]. Note that in this fitting procedure only the asymptotic forms of




FIG. 1: Twist-2 light-cone wave functions for (a) the pion $\tilde{\mathcal{P}}_{2;\pi}$ and (b) the kaon $\tilde{\mathcal{P}}_{2;K}$ (solid lines), along with the wave functions corresponding to the respective asymptotic DAs (dashed lines). The DAs are defined at the scale $\mu_0 = 1$ GeV.

the twist-2 and twist-3 DAs (Eq. (9) & Eq. (20)) were used in the pion wave functions. The running behavior $\chi_{3M}(\mu)$ is then later introduced while calculating the form factors whose RG behavior is assumed to be the same as that of μ_M (see, Appendix A). In other words, this amounts to the replacement $\mu_M^2 \rightarrow \chi_{3M}(1/b_1) \chi_{3M}(1/b_2)$ in Eq. (46). Also, as the bulk of the world pion data is concentrated in the very low energy region where the usual running coupling rapidly diverges, we in addition use an analytic prescription for the QCD running coupling to prove our results. The analytic scheme was suggested originally in Ref. [90] for calculating the pion form factor and further developed in Refs. [60, 61] for NLO calculations. Here, the central idea is the removal of the explicit Landau singularity present in perturbation theory rendering the coupling constant IR stable and reducing the IR sensitivity of perturbatively calculated hadronic observables. The scheme is also known to display higher loop stability. Now, the usual two-loop running coupling Eq. (31) in standard pQCD can be approximately expressed via the *Lambert* W_{-1} function

$$\frac{\alpha_s(\mu^2)}{\pi} = -\frac{\beta_0}{\beta_1} \left[1 + W_{-1} \left(-\frac{\beta_0^2}{\beta_1 e} \left(\frac{\Lambda_{\text{QCD}}^2}{\mu^2} \right) \right) \right]^{-1}. \quad (63)$$

The extension of the above formula in analytic perturbation theory is too complicated to be evaluated exactly and instead there is an alternate approximate expression in the $\overline{\text{MS}}$




FIG. 2: 2-particle twist-3 light-cone wave functions for the pion (a) $\tilde{\mathcal{P}}_{3;\pi}^p$ & (b) $\tilde{\mathcal{P}}_{3;\pi}^\sigma$ and for the kaon (c) $\tilde{\mathcal{P}}_{3;K}^p$ & (d) $\tilde{\mathcal{P}}_{3;K}^\sigma$ with G-parity even terms (solid lines), along with the wave functions corresponding to the respective asymptotic DAs (long dashed lines). For the kaon, the twist-3 wave functions including G-parity odd terms are also shown (dotted lines). The DAs are defined at the scale $\mu_0 = 1$ GeV.

scheme, as suggested in Ref. [89] :

$$\frac{\alpha_s^{\text{an,approx}}(\mu^2)}{\pi} = \frac{1}{\beta_0} \left[\frac{1}{l} + \frac{1}{1 - \exp(l)} \right] ; \quad (64)$$

$\pi^\pm$	At $\mu_0 = 1$ GeV	$K^\pm$	At $\mu_0 = 1$ GeV	units
-	-	$m_{u,d}$	5.6 ± 1.6 [37]	MeV
-	-	m_s	137 ± 27 [37]	MeV
$\mathcal{M}_{u,d}$	0.33	$\mathcal{M}_{u,d}$	0.33	GeV
-	-	$\mathcal{M}_s$	0.45	GeV
m_π	139	m_K	493	MeV
a_1^π	0	a_1^K	0.06 ± 0.03 [37]	-
a_2^π	0.25 ± 0.15 [37]	a_2^K	0.25 ± 0.15 [37]	-
f_π	131	f_K	$1.22 f_\pi $ [24]	MeV
$f_{3\pi}$	0.0045 ± 0.0015 [37]	f_{3K}	0.0045 ± 0.0015 [37]	GeV ²
$\omega_{3\pi}$	-1.5 ± 0.7 [37]	ω_{3K}	-1.2 ± 0.7 [37]	-
$\lambda_{3\pi}$	0	λ_{3K}	1.6 ± 0.4 [37]	-

TABLE I: Various input hadronic parameters for twist-2 and twist-3 light-cone wave functions at $\mu_0 = 1$ GeV.

$$l = \ln \left(\frac{\mu^2}{\Lambda_{\text{an}}^2} \right) + \frac{\beta_1}{\beta_0^2} \ln \sqrt{\ln^2 \left(\frac{\mu^2}{\Lambda_{\text{an}}^2} \right) + 4\pi^2}, \quad (65)$$

where Λ_{an} in the analytic scheme is the analog of Λ_{QCD} in the usual perturbation theory and is chosen to be around 0.4 GeV for $N_f = 3$. We use this formula for the analytic coupling in our calculations. The simple one-parameter fitting of the total form factor to the experimental data gives the best fit values of 1.2 GeV and 1.4 GeV for the usual and analytic QCD coupling schemes, respectively. Here, we choose the average value $\chi_{3M} = 1.3$ GeV for both the schemes, and generously consider the resulting difference from the phenomenological value of 1.7 GeV to contribute to the theoretical error, i.e., ± 0.4 GeV. Note that the fitting takes into account the individual error-bars of the data points.

Finally, following Ref. [95] the Sudakov suppression factor $\exp(-s(XQ, 1/b))$ is set to unity for small transverse separation “ b ” between the valence quarks, i.e., whenever $b < \sqrt{2}/(XQ)$. Also, to avoid probing into certain kinematic regions where $\exp(-S)$ may become greater than unity causing an enhancement instead of a suppression, $\exp(-S)$ is set to 1 for $S < 0$.

$\pi^\pm$	G (even)	$K^\pm$	G (even)	G (even + odd)	units
$A_{2;\pi}$	1.69 (1.66) _{as}	$A_{2;K}$	2.06 (2.07) _{as}	2.06 (2.07) _{as}	-
$A_{3;\pi}^p$	3.76 (3.59) _{as}	$A_{3;K}^p$	4.40 (4.56) _{as}	4.35 (4.56) _{as}	-
$A_{3;\pi}^\sigma$	3.37 (3.33) _{as}	$A_{3;K}^\sigma$	4.16 (4.14) _{as}	4.06 (4.14) _{as}	-
$(\beta_{2;\pi})^2$	0.76 (0.87) _{as}	$(\beta_{2;K})^2$	0.78 (0.89) _{as}	0.78 (0.89) _{as}	GeV ⁻²
$(\beta_{3;\pi}^p)^2$	0.62 (0.74) _{as}	$(\beta_{3;K}^p)^2$	0.70 (0.79) _{as}	0.65 (0.79) _{as}	GeV ⁻²
$(\beta_{3;\pi}^\sigma)^2$	0.81 (0.87) _{as}	$(\beta_{3;K}^\sigma)^2$	0.88 (0.89) _{as}	0.84 (0.89) _{as}	GeV ⁻²
$\chi_{3\pi}^{\text{fit}}$	1.3 ± 0.4	χ_{3K}^{fit}	1.3 ± 0.4	1.3 ± 0.4	GeV

TABLE II: Various determined hadronic parameters for twist-2 and twist-3 light-cone wave functions at $\mu_0 = 1$ GeV. The numbers in the parentheses correspond to values for the asymptotic wave functions.

A. The pion form factor

Using the DAs in Appendix A, we evaluated the total electromagnetic form factor for the pion Eq. (54) using both the usual two-loop QCD running coupling Eq. (31) and the analytical prescription Eq. (64). Fig. 3 shows our results for the total form factor, along with the experimental “world-data” [1, 2, 3, 4, 5, 6, 7, 8] for the pion. The plots correspond to $\Lambda_{\text{QCD}} = 0.2$ GeV and $\Lambda_{\text{an}} = 0.4$ GeV, respectively. It appears that the full twist-3 calculations improve the agreement with experimental data to a much better extent at intermediate energies down to around 1-2 GeV² than for the twist-2 case. Note that in the usual perturbative scheme, as $Q^2 \rightarrow 0$ the total form factor becomes very unpredictable and starts oscillating between large values although $F_\pi(Q^2 = 0) = 1$, satisfying the Ward identity. This clearly signals the breakdown of perturbation theory at such small momentum transfers.

To study the contributions of endpoint effects and to distinguish individual soft and hard contributions, it is more useful to study the variation of the scaled pion form factor $Q^2 F_\pi$ with Q^2 . In Fig. 4, we show the individual contributions of the twist-2 and twist-3 power correction to the scaled hard pion form factor over a wide range of momentum transfers for the usual QCD coupling. Clearly, the twist-3 contributions are seen to be significantly larger than the leading twist-2 counterparts at low momentum transfers, supporting the claims




FIG. 3: The total electromagnetic pion form factor at the twist-2 level (Soft+Twist-2), denoted by the dashed lines and at the twist-3 level (Soft+Twist-2+Twist-3), denoted by the solid lines, with (a) the usual QCD running coupling and (b) the analytical QCD running coupling. The world pion data are taken from Refs. [1, 2, 3, 4, 5, 6, 7, 8].

made in [25, 27, 28, 57, 62, 64]. In fact, it is interesting to see the endpoint enhancement in the twist-3 amplitudes much more explicitly if one rather considered only the collinear DAs to calculate the form factors in the usual perturbation theory, without considering the full transverse momentum dependence (e.g., the BHL ansatz) in the meson wave functions, as originally done in Ref. [95]. In other words, one simply makes the replacement $\tilde{\mathcal{P}}_M(x) \rightarrow \phi_M(x)$ in calculating the hard form factor. The inclusion of the transverse momenta and constituent quark masses in the wave function provides a natural cut-off for the soft and endpoint enhancements. Similar behavior can also be observed in the analytic case, although we have not displayed the enhancement which is less drastic. These facts suggest that the modified collinear factorization scheme, including explicit transverse degrees of freedom with Sudakov suppression, which works well for the twist-2 case is not very effective at the




FIG. 4: Twist-2 (long dashed lines) and Twist-3 (solid lines) corrections to the scaled hard pion form factor with the usual QCD running coupling. Plots (a) & (b) are obtained using BHL ansatz, while (c) & (d) are obtained with $\tilde{\mathcal{P}}_M(x) \rightarrow \phi_M(x)$. Also, the plots (a) & (c) do not include threshold resummation in the hard contributions, which are included in (b) & (d). The soft corrections (short dashed lines) are also shown.

twist-3 level in shielding such artificial non-perturbative enhancements at low momenta. To improve this situation, especially for the results obtained in usual perturbative scheme, we use threshold resummation which along with Sudakov suppression provide large damping of

the endpoint effects in the twist-3 amplitude. The twist-2 part on the other hand remains mostly unaltered, if not slightly enhanced due to the threshold resummation, especially in the low-energy region. Note that in this respect the use of threshold resummation in the analytic scheme is somewhat redundant and has little effect on both the twist corrections. Finally, as expected, one observes that the twist-3 corrections fall off rapidly with increasing Q^2 and beyond a certain point fall below the twist-2 corrections. At asymptotically large momentum transfers, only the twist-2 contributions are expected to dominate.

Our final results for the scaled pion form factor are summarized in Fig. 5 and Fig. 6. We use both the usual and the analytic QCD running couplings and compare our results with the available experimental pion data with increasing error bars towards intermediate energies. The individual soft and hard contributions along with the total contribution are shown. Clearly, contrary to the earlier claims made in Ref. [58], the twist-2 hard form factor is far too small in the phenomenologically accessible region to explain the data. One must therefore look for other possibilities like non-perturbative higher twist effects and soft contributions. Interestingly, it is seen that the soft dynamics largely dominate the low-energy region below 10-16 GeV² but rapidly fall off in the asymptotic region. The contributions from the twist-3 corrections turn out to be significantly large in the moderate range of energies below 100 GeV², but eventually the hard twist-2 contributions solely determine the asymptotic trend beyond $Q^2 \approx 100\text{-}150$ GeV².

As evident from the figures, the total scaled pion form factor (solid lines) up to twist-3 corrections displays an obvious improvement of the overall agreement with experimental data compared to the twist-2 scaled form factor (dashed lines). To some extent, it is somewhat surprising to see that in combination with the soft contributions, the modified resummed pQCD with the usual QCD coupling could work so well as low as $Q^2 \approx 0.25$ GeV², far lower than previously envisaged. To this end, we display the results in the analytic scheme to confirm our results. The analytical prescription is known to reduce the scheme and renormalization scale dependence, largely increasing the stability of solutions [61]. Accordingly, there is some confidence in our displayed results. The results obtained in both schemes not only show a striking similarity even at sufficiently low momentum transfers, but also show a good agreement with the available pion form factor data. However, whether or not such an agreement is merely accidental is matter of debate. There may still be substantial sub-leading contributions e.g., from a full NLO calculation in the strong coupling including




FIG. 5: Scaled total electromagnetic form factor for the pion with (a) the usual QCD running coupling and (b) the analytical running coupling in the low and intermediate energy regime. The solid line represents the full twist-3 result (Soft+Twist-2+Twist-3), the long dashed lines represent the twist-2 result (Soft+Twist-2) and the soft corrections are indicated by the short dashed lines. The world pion data are taken from Refs. [1, 2, 3, 4, 5, 6, 7, 8].




FIG. 6: Scaled total electromagnetic form factor for the pion with (a) the usual QCD running coupling and (b) the analytical running coupling over a wider range of the intermediate energies. The solid line represents the full twist-3 result (Soft+Twist-2 +Twist-3), the long dashed lines represent the full twist-2 result (Soft+Twist-2) and the soft corrections are indicated by the short dashed lines. The world pion data are taken from Refs. [1, 2, 3, 4, 5, 6, 7, 8].

subleading twists and intrinsic transverse momenta for the hard scattering kernel and the DAs, or from higher order Fock states and helicity components in the light-cone DAs. Note that a NLO calculation in the strong QCD coupling constant was done in Ref. [60] where the corrections to the twist-2 pion form factor was found to be quite large. A full NLO calculation for subleading twists, including also the transverse momentum dependence, is however, still missing. In Ref. [63], contributions of higher helicity states were found to lower the total pion form factor significantly. Hence, without systematically taking all of these effects in account, which is beyond the scope of the paper, no definitive statement can be made as to how well our results agree with the data. Moreover, the available data itself has very low statistics at intermediate energy and is plagued with large uncertainties. It is therefore difficult to give a proper theoretical error estimate of our results, when the asymptotic formalism is itself largely unreliable in the region of our interest. What we have done in this paper is a combination of model and pQCD calculation. It may thus be worth using the estimates for the ranges over which the input parameters of the DAs, namely $\mu_\pi(\chi_{3\pi})$, $f_{3\pi}$, $\omega_{3\pi}$ and a_2^π vary (given in the Tables I & II), in computing our theoretical error. In addition, we




FIG. 7: Theoretical 1σ -error for the scaled total pion form factor due to the variation of the input parameters for the DAs ($\chi_{3\pi}$, $f_{3\pi}$, $\omega_{3\pi}$ and a_2^π) and $\Lambda_{\text{QCD,an}}$. The lines correspond to the mean values of the parameters using the usual (dotted line) and the analytic (dashed line) QCD coupling schemes, respectively. The error bars for the experimental data points [1, 2, 3, 4, 5, 6, 7, 8] are also shown.

allow a variation of ± 0.05 GeV for both Λ_{QCD} and Λ_{an} . We used a Monte Carlo technique to generate a gaussian “ 1σ ” spread of the scaled form factor for various Q^2 values about a central mean. The error estimate displayed in Fig. 7 shows our maximum theoretical error to be about 10 percent for the usual QCD coupling and somewhat less for the analytic coupling.




FIG. 8: Scaled total electromagnetic form factor for the kaon with the usual QCD running coupling (upper plot) and the analytical running coupling (lower plot) for intermediate energies. The solid lines represents the full twist-3 result (Soft+Twist-2 +Twist-3), the long dashed lines represent the twist-2 result (Soft+Twist-2) and the soft corrections are indicated by the short dashed lines. For the full twist-3 case, both the results i.e., with and without including the G-parity-breaking terms in the DAs are shown.

B. The kaon form factor

We conclude the section on the numerical analysis by displaying our predictions for the kaon form factor, applying the same techniques as for the case of the pion form factor. The twist-2 and twist-3 light-cone wave functions for the kaon has the general form in the transverse b -space given by

$$\begin{aligned} \tilde{\mathcal{P}}_K(x, b, 1/b, \mathcal{M}_{u,d,s}) = & A_K \phi_K(x, 1/b) \exp \left[-\frac{b^2 x(1-x)}{4\beta_K^2} \right] \\ & \times \exp \left[-\beta_K^2 \left(\frac{\mathcal{M}_s^2}{x} + \frac{\mathcal{M}_{u,d}^2}{1-x} \right) \right], \end{aligned} \quad (66)$$

where we used the twist-3 chiral enhancement parameter $\chi_{3K} = 1.3$ GeV and the experimental estimate for the kaon decay constant $f_K \approx 1.22 f_\pi$ [24]. Here, we also take $\Lambda_{\text{QCD}} = 0.2$ GeV and $\Lambda_{\text{an}} = 0.4$ GeV for the respective running couplings in the $\overline{MS}$ scheme. The results are summarized in Fig. 8 for intermediate energies. The solid line represents the total scaled form factor in each case. Here, our results for the kaon form factor must be considered preliminary. Due to the complete absence of experimental data at intermediate energies we are unable to make any meaningful phenomenological comparison. The presently available kaon data has very poor statistics and has hardly been measured above 0.2 GeV^2 . Hence, we do not show the experimental data points in the form factor plots. With the availability of better quality data in future there could be plenty of room for further improvements, for instance, extension of the above results to include a full NLO calculation for subleading twists or higher helicity and Fock state contributions.

V. DISCUSSION AND CONCLUSION

For the past two decades the electromagnetic meson form factors have been the subject of intensive theoretical and experimental scrutiny and yet there is still to be an universally accepted framework for their description. Presently, reliable experimental data are available only for the pions which are entirely concentrated at very low energies with very poor statistics at intermediate energies. The low energy part of the data is best explained by the standard VMD model, showing no apparent trace of pQCD behavior, which is expected only at very high energies. Very many attempts have been made to predict the onset of the perturbative behavior for the pion form factor. The modified or resummed valence pQCD

with factorization appears to show some attractive features to enable pQCD calculations to be valid in a self-consistent way even at very moderate energies. Whether this is true, will only be confirmed when data with better statistics at higher momentum transfers will become available in the future. At the same time, the onset of the perturbative behavior being very slow, it is still unclear whether the leading order perturbative calculations could be expected to be precise even at the highest accessible energies. In this paper, with the help of (a) the double-humped type DAs and (b) the modified transverse (k_T) factorization scheme, incorporating both Sudakov suppression and threshold resummation, we got rid of non-perturbative end-point enhancements. This enlarges the scope of applicability of resummed pQCD independent of the coupling scheme to a much wider range of intermediate energies, if not down to a few GeVs, as demonstrated in this paper. By a simple adjustment of only the chiral enhancement parameter $\chi_{3\pi}$, a good agreement with the experimental data was obtained.

As for the scaled pion form factor, we found that the leading order pQCD contributions potentially undermine the agreement with the available low energy data and are only trustworthy in the hard energy regime: even from a very conservative point of view, Q^2 should be bigger than 4 GeV². At low momentum transfers, the non-perturbative contributions dominate, being larger than the hard (twist-2) contributions at least by a factor of two. In fact below 4 GeV², 60-70% of the available data is already accounted for by the soft contributions. In addition, we also needed the twist-3 power corrections to explain the remaining discrepancy. However, at larger energies (say, $Q^2 > 50$ -100 GeV²), both the soft and the twist-3 contributions rapidly fall off and eventually the twist-2 form factor dominates asymptotically. Similar conclusions, albeit being preliminary, are drawn for the kaon form factor although it seems that the onset of the perturbative behavior occurs at slightly larger momentum transfers than for the case of the pion. Of course, as we mentioned earlier, it still remains to be investigated about the nature of the contributions that may arise from a full systematic NLO calculation with subleading twists and intrinsic transverse momenta, or from the inclusion of higher helicity and Fock states. With the availability of better quality of data in future such analyses may be necessary to make definite conclusions.

In summary, although the quality of present experimental form factor data does not allow a definitive conclusion, one can expect that the non-perturbative soft contributions and higher twist power corrections to the form factors play an important role at phe-

nomenologically accessible momentum transfers. Thus, more work is needed to be done on both the theoretical and experimental sides to obtain more conclusive results and push the frontiers of our knowledge on confinement dynamics through the study of higher order and non-perturbative contribution to exclusive processes.

Acknowledgments: The authors would like to thank Alexander Bakulev, Pankaj Jain, Alexander Lenz, Seregy Mikhailov, Sabyasachi Mishra, Ingo Sick, Ritesh Singh, Nikolaos Stefanis and Xing-Gang Wu for interesting discussions and comments. One of the authors, U.R. would like to convey special thanks to Prof. Dirk Trautmann and Prof. Friedel Thielemann for their hospitality and financial support at the University of Basel. This work was supported by the Swiss National Science Foundation (SNSF) under the contract number 200020-111705.

APPENDIX A: 2-PARTICLE TWIST-3 DISTRIBUTION AMPLITUDES

The twist-3 DAs are obtained by an expansion over conformal spins. At next-to-leading order, the 2-particle DAs $\phi_{3,M}^p$ and $\phi_{3,M}^\sigma$ (including meson-mass corrections that break chiral symmetry at $\mathcal{O}(m_s + m_q)$ in the SU(3) case while preserving G-parity) are given in terms of the Gegenbauer polynomials $C_n^{1/2}$ and $C_n^{3/2}$, respectively [34, 35] as

$$\begin{aligned}\phi_{3,M}^p(x, \mu^2) &= \frac{f_M}{4\sqrt{2N_c}} \left\{ 1 + \left(30\eta_{3M}(\mu^2) - \frac{5}{2}\rho_M^2(\mu^2) \right) C_2^{1/2}(\xi) \right. \\ &\quad \left. + \left(-3\eta_{3M}(\mu^2)\omega_{3M}(\mu^2) - \frac{27}{20}\rho_M^2(\mu^2) - \frac{81}{10}\rho_M^2(\mu^2)a_2^M(\mu^2) \right) C_4^{1/2}(\xi) \right\}, \\ \phi_{3,M}^\sigma(x, \mu^2) &= \frac{3f_M}{2\sqrt{2N_c}} x(1-x) \left\{ 1 + \left(5\eta_{3M}(\mu^2) - \frac{1}{2}\eta_{3M}(\mu^2)\omega_{3M}(\mu^2) \right. \right. \\ &\quad \left. \left. - \frac{7}{20}\rho_M^2(\mu^2) - \frac{3}{5}\rho_M^2(\mu^2)a_2^M(\mu^2) \right) C_2^{3/2}(\xi) \right\}\end{aligned}\quad (\text{A1})$$

with

$$\eta_{3M} = \frac{f_{3M}}{f_M} \frac{1}{\mu_M} \quad ; \quad \rho_M = \frac{m_M}{\mu_M} \quad ; \quad M = \pi^\pm, K^\pm \quad (\text{A2})$$

where the non-perturbative parameter f_{3M} and ω_{3M} , respectively are defined by the following matrix elements of local twist-3 operators:

$$\langle 0 | \bar{q}_{f_1} \sigma_{\mu\nu} \gamma_5 g_s G_{\alpha\beta} q_{f_2} | M(P) \rangle = i f_{3M} (P_\alpha P_\mu g_{\nu\beta} - P_\alpha P_\nu g_{\mu\beta} - P_\beta P_\mu g_{\nu\alpha} + P_\beta P_\nu g_{\alpha\mu}),$$

$$\begin{aligned}\langle 0 | \bar{q}_{f_1} \sigma_{\mu\lambda} \gamma_5 [iD_\beta, g_s G_{\alpha\lambda}] q_{f_2} - (3/7) i \partial_\beta \bar{q}_{f_1} \sigma_{\mu\lambda} \gamma_5 g_s G_{\alpha\lambda} q_{f_2} | M(P) \rangle \\ = \frac{3}{14} i f_{3M} P_\alpha P_\beta P_\mu \omega_{3M},\end{aligned}\quad (\text{A3})$$

where $G_{\alpha\beta}$ is the gluon field tensor. The LO scale dependence of various twist-3 parameters are given by

$$\begin{aligned}\rho_M(\mu^2) &= L^{\gamma_{3;q\bar{q}}^{(0)}/\beta_0} \rho_M(\mu_0^2); \quad \gamma_{3;q\bar{q}}^{(0)} = 1, \\ \mu_M(\mu^2) &= L^{\gamma_{3;\mu}^{(0)}/\beta_0} \mu_M(\mu_0^2); \quad \gamma_{3;\mu}^{(0)} = -\gamma_{3;q\bar{q}}^{(0)} = -1, \\ \eta_{3M}(\mu^2) &= L^{\gamma_{3;\eta}^{(0)}/\beta_0} \eta_{3M}(\mu_0^2); \quad \gamma_{3;\eta}^{(0)} = \frac{4}{3}C_F + \frac{1}{4}C_A, \\ \omega_{3M}(\mu^2) &= L^{\gamma_{3;\omega}^{(0)}/\beta_0} \omega_{3M}(\mu_0^2); \quad \gamma_{3;\omega}^{(0)} = -\frac{7}{24}C_F + \frac{7}{12}C_A, \\ a_2^M(\mu^2) &= L^{\gamma_2^{(0)}/\beta_0} a_2^M(\mu_0^2); \quad \gamma_2^{(0)} = \frac{25}{24}C_F,\end{aligned}\quad (\text{A4})$$

where $L = \alpha_s(\mu^2)/\alpha_s(\mu_0^2)$, $\mathcal{C}_F = (N_c^2 - 1)/2N_c$, $\mathcal{C}_A = N_c$ and $\mu_0 \approx 1$ GeV.

We have also considered the 2-particle twist-3 kaon DAs $\phi_{3,K}^p$ and $\phi_{3,K}^\sigma$, as given in [37], which not only include a complete set of meson-mass corrections but also G-parity-breaking terms of $\mathcal{O}(m_s - m_q)$:

$$\begin{aligned} \phi_{3,K}^p(x, \mu^2) = & \frac{f_K}{4\sqrt{2N_c}} \left\{ 1 + 3\rho_+^K(1 + 6a_2^K) - 9\rho_-^K a_1^K + \left[\frac{27}{2} \rho_+^K a_1^K \right. \right. \\ & \left. \left. - \rho_-^K \left(\frac{3}{2} + 27a_2^K \right) \right] C_1^{1/2}(\xi) + \left(30\eta_{3K} + 15\rho_+^K a_2^K - 3\rho_-^K a_1^K \right) C_2^{1/2}(\xi) \right. \\ & \left. + \left(10\eta_{3K}\lambda_{3K} - \frac{9}{2}\rho_-^K a_2^K \right) C_3^{1/2}(\xi) - 3\eta_{3K}\omega_{3K} C_4^{1/2}(\xi) + \frac{3}{2} (\rho_+^K + \rho_-^K) \right. \\ & \left. \times (1 - 3a_1^K + 6a_2^K) \ln x + \frac{3}{2} (\rho_+^K - \rho_-^K) (1 + 3a_1^K + 6a_2^K) \ln(1 - x) \right\}, \end{aligned} \quad (\text{A5})$$

$$\begin{aligned} \phi_{3,K}^\sigma(x, \mu^2) = & \frac{3f_K}{2\sqrt{2N_c}} x(1 - x) \left\{ 1 + \frac{3}{2} \rho_+^K + 15\rho_+^K a_2^K - \frac{15}{2} \rho_-^K a_1^K \right. \\ & \left. + \left(3\rho_+^K a_1^K - \frac{15}{2} \rho_-^K a_2^K \right) C_1^{3/2}(\xi) + \left(5\eta_{3K} - \frac{1}{2} \eta_{3K}\omega_{3K} + \frac{3}{2} \rho_+^K a_2^K \right) C_2^{3/2}(\xi) \right. \\ & \left. + \eta_{3K}\lambda_{3K} C_3^{3/2}(\xi) + \frac{3}{2} (\rho_+^K + \rho_-^K) (1 - 3a_1^K + 6a_2^K) \ln x \right. \\ & \left. + \frac{3}{2} (\rho_+^K - \rho_-^K) (1 + 3a_1^K + 6a_2^K) \ln(1 - x) \right\} \left(\frac{1}{1 - \rho_+} \right) \end{aligned} \quad (\text{A6})$$

with

$$\eta_{3K} = \frac{f_{3K}}{f_K} \frac{1}{\mu_K}, \quad \rho_+^K = \frac{(m_s + m_q)^2}{m_K^2} \quad \text{and} \quad \rho_-^K = \frac{m_s^2 - m_q^2}{m_K^2}. \quad (\text{A7})$$

Note that the expression for $\phi_{3,K}^\sigma(x, \mu^2)$ is normalized to unity with an extra factor of $1/(1 - \rho_+)$, compared to that given in [37]. The non-perturbative parameters f_{3K} , ω_{3K} and λ_{3K} are defined (e.g., K^-) by

$$\langle 0 | \bar{u} \sigma_{\mu\nu} \gamma_5 g_s G_{\alpha\beta} s | K^-(P) \rangle = i f_{3K} (P_\alpha P_\nu g_{\nu\beta} - P_\alpha P_\nu g_{\mu\beta} - P_\beta P_\mu g_{\nu\alpha} + P_\beta P_\nu g_{\alpha\mu}),$$

$$\langle 0 | \bar{u} \sigma_{\mu\lambda} \gamma_5 [iD_\beta, g_s G_{\alpha\lambda}] s - (3/7) i \partial_\beta \bar{u} \sigma_{\mu\lambda} \gamma_5 g_s G_{\alpha\lambda} s | K^-(P) \rangle = \frac{3}{14} i f_{3K} P_\alpha P_\beta P_\mu \omega_{3K},$$

$$\langle 0 | \bar{u} i \overleftrightarrow{D}_\beta \sigma_{\mu\lambda} \gamma_5 g_s G_{\alpha\lambda} s - \bar{u} \sigma_{\mu\lambda} \gamma_5 g_s G_{\alpha\lambda} i \overrightarrow{D}_\beta s | K^-(P) \rangle = \frac{1}{7} i f_{3K} P_\alpha P_\beta P_\mu \lambda_{3K}. \quad (\text{A8})$$

where in the chiral limit the renormalization group equations at LO give,

$$\begin{aligned}
\mu_K(\mu^2) &= L^{\gamma_{3;\mu}^{(0)}/\beta_0} \mu_K(\mu_0^2); \quad \gamma_{3;\mu}^{(0)} = -\gamma_{3;q\bar{q}}^{(0)} = -1, \\
\rho_+^K(\mu^2) &= L^{\gamma_{3;\rho^+}^{(0)}/\beta_0} \rho_+^K(\mu_0^2); \quad \gamma_{3;\rho^+}^{(0)} = 2\gamma_{3;q\bar{q}}^{(0)} = 2, \\
\rho_-^K(\mu^2) &= L^{\gamma_{3;\rho^-}^{(0)}/\beta_0} \rho_-^K(\mu_0^2); \quad \gamma_{3;\rho^-}^{(0)} = 2\gamma_{3;q\bar{q}}^{(0)} = 2, \\
f_{3K}(\mu^2) &= L^{\gamma_{3;f}^{(0)}/\beta_0} f_{3K}(\mu_0^2); \quad \gamma_{3;f}^{(0)} = \frac{7}{12} \mathcal{C}_F + \frac{1}{4} \mathcal{C}_A, \\
\omega_{3K}(\mu^2) &= L^{\gamma_{3;\omega}^{(0)}/\beta_0} \omega_{3K}(\mu_0^2); \quad \gamma_{3;\omega}^{(0)} = -\frac{7}{24} \mathcal{C}_F + \frac{7}{12} \mathcal{C}_A, \\
\lambda_{3K}(\mu^2) &= L^{\gamma_{3;\lambda}^{(0)}/\beta_0} \lambda_{3K}(\mu_0^2); \quad \gamma_{3;\lambda}^{(0)} = \frac{19}{48} \mathcal{C}_F, \\
a_1^K(\mu^2) &= L^{\gamma_1^{(0)}/\beta_0} a_1^K(\mu_0^2); \quad \gamma_1^{(0)} = \frac{2}{3} \mathcal{C}_F, \\
a_2^K(\mu^2) &= L^{\gamma_2^{(0)}/\beta_0} a_2^K(\mu_0^2); \quad \gamma_2^{(0)} = \frac{25}{24} \mathcal{C}_F,
\end{aligned} \tag{A9}$$

But the strange quark being massive, there is operator mixing of the ones in Eq. (A8) with that of twist-2 operators and the resulting LO renormalization group equations give the following scale dependence of the various twist-3 parameters:

$$\begin{aligned}
f_{3K}(\mu^2) &= L^{55/(36\beta_0)} f_{3K}(\mu_0^2) + \frac{2}{19} (L^{1/\beta_0} - L^{55/(36\beta_0)}) [f_K m_s](\mu_0^2) \\
&\quad + \frac{6}{65} (L^{55/(36\beta_0)} - L^{17/(9\beta_0)}) [f_K m_s a_1^K](\mu_0^2), \\
[f_{3K} \omega_{3K}](\mu^2) &= L^{26/(9\beta_0)} [f_{3K} \omega_{3K}](\mu_0^2) + \frac{1}{170} (L^{1/\beta_0} - L^{26/(9\beta_0)}) [f_K m_s](\mu_0^2) \\
&\quad + \frac{1}{10} (L^{17/(9\beta_0)} - L^{26/(9\beta_0)}) [f_K m_s a_1^K](\mu_0^2) \\
&\quad + \frac{2}{15} (L^{43/(18\beta_0)} - L^{26/(9\beta_0)}) [f_K m_s a_2^K](\mu_0^2), \\
[f_{3K} \lambda_{3K}](\mu^2) &= L^{37/(18\beta_0)} [f_{3K} \lambda_{3K}](\mu_0^2) - \frac{14}{67} (L^{1/\beta_0} - L^{37/(18\beta_0)}) [f_K m_s](\mu_0^2) \\
&\quad + \frac{14}{5} (L^{17/(9\beta_0)} - L^{37/(18\beta_0)}) [f_K m_s a_1^K](\mu_0^2) \\
&\quad - \frac{4}{11} (L^{43/(18\beta_0)} - L^{37/(18\beta_0)}) [f_K m_s a_2^K](\mu_0^2).
\end{aligned} \tag{A10}$$

Finally, we present the various Gegenbauer polynomials used in the above formulae:

$$\begin{aligned}
C_1^{1/2}(\xi) &= \xi, \\
C_2^{1/2}(\xi) &= \frac{1}{2}(3\xi^2 - 1), \\
C_3^{1/2}(\xi) &= \frac{1}{2}\xi(5\xi^2 - 3), \\
C_4^{1/2}(\xi) &= \frac{1}{8}(35\xi^4 - 30\xi^2 + 3), \\
C_0^{3/2}(\xi) &= 1, \\
C_1^{3/2}(\xi) &= 3\xi, \\
C_2^{3/2}(\xi) &= \frac{3}{2}(5\xi^2 - 1), \\
C_3^{3/2}(\xi) &= \frac{5}{2}\xi(7\xi^2 - 3).
\end{aligned} \tag{A11}$$

APPENDIX B: CALCULATION OF THE SUDAKOV EXPONENT

The full expression of the Sudakov suppression factor $S(x, y, b_1, b_2, Q)$ is given by,

$$\begin{aligned}
S(x, y, b_1, b_2, Q) &= s(xQ, b_1) + s(yQ, b_2) + s((1-x)Q, b_1) + s((1-y)Q, b_2) \\
&\quad - \frac{1}{\beta_0} \ln \left(\frac{\hat{t}}{-\hat{b}_1} \right) - \frac{1}{\beta_0} \ln \left(\frac{\hat{t}}{-\hat{b}_2} \right) \\
&\quad + \frac{\beta_1}{\beta_0^3} \left[\frac{1 + \ln(-2\hat{b}_1)}{-2\hat{b}_1} - \frac{1 + \ln(2\hat{t})}{2\hat{t}} \right] \\
&\quad + \frac{\beta_1}{\beta_0^3} \left[\frac{1 + \ln(-2\hat{b}_2)}{-2\hat{b}_2} - \frac{1 + \ln(2\hat{t})}{2\hat{t}} \right],
\end{aligned} \tag{B1}$$

where

$$\begin{aligned}
s(XQ, 1/b) = & \frac{\mathcal{A}^{(1)}}{2\beta_0} \hat{q} \ln \left(\frac{\hat{q}}{-\hat{b}} \right) + \frac{\mathcal{A}^{(2)}}{4\beta_0^2} \left(\frac{\hat{q}}{-\hat{b}} - 1 \right) - \frac{\mathcal{A}^{(1)}}{2\beta_0} (\hat{b} + \hat{q}) \\
& - \left[\frac{4\mathcal{A}^{(1)}\beta_1}{16\beta_0^3} \hat{q} + \frac{2\mathcal{A}^{(1)}\beta_1}{16\beta_0^3} \ln \left(\frac{1}{2} e^{2\gamma_E - 1} \right) \right] \left[\frac{1 + \ln(-2\hat{b})}{-\hat{b}} - \frac{1 + \ln(2\hat{q})}{\hat{q}} \right] \\
& - \left[\frac{\mathcal{A}^{(2)}}{4\beta_0^2} - \frac{\mathcal{A}^{(1)}}{4\beta_0} \ln \left(\frac{1}{2} e^{2\gamma_E - 1} \right) \right] \ln \left(\frac{\hat{q}}{-\hat{b}} \right) - \frac{4\mathcal{A}^{(1)}\beta_1}{32\beta_0^3} [\ln^2(-2\hat{b}) - \ln^2(2\hat{q})] \\
& + \frac{2\mathcal{A}^{(2)}\beta_1}{8\beta_0^4} \left[\frac{1 + \ln(-2\hat{b})}{-\hat{b}} - \frac{1 + \ln(2\hat{q})}{\hat{q}} \right] \\
& - \frac{2\mathcal{A}^{(2)}\beta_1}{8\beta_0^4} \hat{q} \left[\frac{1 + 2\ln(-2\hat{b})}{(-2\hat{b})^2} - \frac{1 + 2\ln(2\hat{q})}{(2\hat{q})^2} \right] \\
& - \frac{4\mathcal{A}^{(2)}\beta_1^2}{64\beta_0^6} \left[\frac{1 + 2\ln(-2\hat{b}) + 2\ln^2(-2\hat{b})}{(-2\hat{b})^2} - \frac{1 + 2\ln(2\hat{q}) + 2\ln^2(2\hat{q})}{(2\hat{q})^2} \right] \\
& + \frac{4\mathcal{A}^{(2)}\beta_1^2}{8\beta_0^6} \hat{q} \left[\frac{\frac{2}{27} + \frac{2}{9}\ln(-2\hat{b}) + \frac{1}{3}\ln^2(-2\hat{b})}{(-2\hat{b})^3} - \frac{\frac{2}{27} + \frac{2}{9}\ln(2\hat{q}) + \frac{1}{3}\ln^2(2\hat{q})}{(2\hat{q})^3} \right].
\end{aligned} \tag{B2}$$

In the above formulae,

$$\begin{aligned}
\hat{t} &= \ln \left(\frac{t}{\Lambda_{\text{QCD}}} \right); \quad t = \max(\sqrt{xy} Q, 1/b_1, 1/b_2), \\
\hat{b} &= \ln(b \Lambda_{\text{QCD}}), \\
\hat{q} &= \ln \left[\frac{XQ}{\sqrt{2}\Lambda_{\text{QCD}}} \right]; \quad X = x, y, (1-x) \text{ or } (1-y), \\
\mathcal{A}^{(1)} &= \mathcal{C}_F = \frac{4}{3}, \\
\mathcal{A}^{(2)} &= \left(\frac{67}{27} - \frac{\pi^2}{9} \right) N_c - \frac{10}{27} N_f + \frac{8}{3} \beta_0 \ln \left(\frac{e^{\gamma_E}}{2} \right).
\end{aligned} \tag{B3}$$

-
- [1] C. N. Brown et al., Phys. Rev. D **8** (1973) 92.
 - [2] C. J. Babek et al., Phys. Rev. D **17** (1978) 1693.
 - [3] H. Ackermann et al., Nucl. Phys. B **137** (1978) 294.
 - [4] P. Brauel et al., Z. Phys. C **3** (1979) 101.
 - [5] S. R. Amendolia et al., Phys. Lett. B **178** (1986) 435; Phys. Lett. B **277** (1986) 168.
 - [6] J. Volmer, Ph.D. thesis, Vrije Universiteit, Amsterdam, 2000 (unpublished); Phys. Rev. Lett. **86** (2001) 1713.

- [7] T. Horn et al., Phys. Rev. Lett. **97** (2006) 192001.
- [8] V. Tadevosyan et al., Phys. Rev. C **75** (2007) 055205.
- [9] B. Zeidman et al., CEBAF Experiment E91-016/1996.
- [10] R. Mohring et al., Phys. Rev. Lett. **81** (1998) 1805.
- [11] M. Vanderhaeghen, M. Guidal and J.-M. Laget, Phys. Rev. C **57** (1998) 1454.
- [12] C. Bennhold and T. Mart, Phys. Rev. C **61** (1999) 012201.
- [13] R. A. Williams, Phys. Rev. C **46** (1992) 1617.
- [14] E. B. Dally et al., Phys. Rev. Lett. **45** (1980) 232.
- [15] K. H. Glander et al., Eur. Phys. J. A **19** (2004) 251.
- [16] J. W. C. McNabb et al., Phys. Rev. C **69** (2004) 042201(R).
- [17] R. G. T. Zegers et al., Phys. Rev. Lett. **91** (2003) 092001.
- [18] B.-W. Xiao and X. Qian, Eur. Phys. J. A **15** (2002) 523 [arXiv:hep-ph/0209138].
- [19] F. P. Pereira et al., Phys. Part. Nucl. **36** (2005) 217 [arXiv:hep-ph/0506032].
- [20] X.-G. Wu, and T. Huang, JHEP 0804:043, 2008 [arXiv:hep-ph/0803.4229].
- [21] J. Bijnens and A. Khodjamirian, Eur. Phys. J. C **26** (2002) 67 [arXiv:hep-ph/0206252].
- [22] J. Botts and G. Sterman, Nucl. Phys. B **325** (1989) 62.
- [23] H.-N. Li and G. Sterman, Nucl. Phys. B **381** (1992) 129.
- [24] V. L. Chernyak and A. R. Zhitnitsky, JETP Lett. **25** (1977) 510; Yad. Fiz. **31** (1980) 1053; Nucl. Phys. B **201** (1982) 492; Sov. J. Nucl. Phys. **38** (1983) 775; Nucl. Phys. B **246** (1984) 52. Phys. Rep. **112** (1984) 173.
- [25] V. L. Chernyak, A. R. Zhitnitsky and I. R. Zhitnitsky Nucl. Phys. B **204** (1982) 477.
- [26] V. L. Chernyak, A. R. Zhitnitsky and V. G. Serbo, JETP Lett. **26** (1977) 594; Sov. J. Nucl. Phys. **31** (1980) 552.
- [27] B. V. Geshkenbeim and M. V. Terentyev, Phys. Lett. B **117** (1982) 243; Sov. J. Nucl. Phys. **39** (1984) 554; Sov. J. Nucl. Phys. **39** (1984) 873.
- [28] C. S. Huang, Commun. Theor. Phys. **2** (1983) 1265.
- [29] M. Gari and N. G. Stefanis, Phys. Lett. B **175** (1986) 462.
- [30] Z. Dziembowski and L. Mankiewicz, Phys. Rev. Lett. **58** (1987) 2175.
- [31] T. Huang and Q. X. Sheng, Z. Phys. C **50** (1991) 139.
- [32] F.-G. Cao, T. Huang and B. Q. Ma, Phys. Rev. D **53** (1996) 6582.
- [33] S. Brodsky and G. F. de Teramond, [arXiv:hep-ph/0804.3562].

- [34] V. M. Braun and I. E. Filyanov, Z. Phys. C **44** (1989) 157.
- [35] P. Ball, JHEP **9901** (1999) 010.
- [36] P. Ball and M. Boglione, Phys. Rev. D **68** (2003) 094006.
- [37] P. Ball, V. M. Braun and A. Lenz, JHEP **0605** (2006) 004 [arXiv:hep-ph/0603063].
- [38] A. Khodjamirian, Th. Mannel and M. Melcher, Phys. Rev. D **68** (2003) 114007; Phys. Rev. D **70** (2004) 094002 [arXiv:hep-ph/0407226].
- [39] C.-R. Ji and F. Amiri, Phys. Rev. D **42** (1990) 3764.
- [40] J. M. Cornwall, Phys. Rev. D **26** (1982) 1453.
- [41] C.-R. Ji, A. F. Sill and R. M. Lombard-Nelson, Phys. Rev. D **36** (1987) 165.
- [42] W. G. Holladay, Phys. Rev. **101** (1956) 1198.
- [43] W. R. Frazer and J. R. Fulco, Phys. Rev. Lett. **2** (1959) 365; Phys. Rev. **117** (1960) 1609.
- [44] J. Sakurai, *Currents and Mesons* (University of Chicago, Chicago, 1969).
- [45] B. Ananthanarayan and S. Ramanan, Eur. Phys. J. C **54** (2008) 461 [arXiv:hep-ph/0801.2023].
- [46] S. J. Brodsky and G. R. Farrar, Phys. Rev. Lett. **31** (1973) 1153; Phys. Rev. D **11** (1975) 1309.
- [47] F. R. Farrar and D. R. Jackson, Phys. Rev. Lett. **43** (1979) 246.
- [48] V. A. Nesterenko and A. V. Radyushkin, Phys. Lett. B **115** (1982) 410.
- [49] V. V. Anisovich, D. I. Melikhov and V. A. Nikonov, Phys. Rev. D **52** (1995) 5295; Phys. Rev. D **55** (1997) 2918.
- [50] A. P. Bakulev, A. V. Radyushkin and N. G. Stefanis, Phys. Rev. D **62** (2000) 113001.
- [51] V. Braguta, W. Lucha and D. Melikhov, Phys. Lett. B **661** (2008) 354 [arXiv:hep-ph/0710.5461].
- [52] C. R. Ji and S. Cotanch, Phys. Rev. D **21** (1990) 2319.
- [53] T. Gousset and B. Pire, Phys. Rev. D **51** (1995) 15; [arXiv:hep-ph/9403293]; Proceedings of the *ELFE Summer School on Confinement physics, Cambridge 1995*, p.111-143 [arXiv:hep-ph/9511274].
- [54] C.-R. Ji, A. Pang and A. Szczepaniak, Phys. Rev. D **52** (1995) 4038.
- [55] J. P. B. C. de Melo et al., Phys. Rev. C **59** (1999) 2278; Nucl. Phys. A **707** (2002) 399
- [56] H.-M. Choi and C.-R. Ji, Phys. Rev. D **74** (2006) 093010 [arXiv:hep-ph/0608148]; [arXiv:hep-ph/0803.2604].
- [57] F.-G. Cao, Y.-B. Dai and C.-S. Huang, Eur. Phys. J. C **11** (1999) 501 [arXiv:hep-ph/9711203].

- [58] C. Coriano, H.-N. Li and C. Savkli, JHEP 9807:008, 1998 [arXiv:hep-ph/9805406].
- [59] V. M. Braun, A. Khodjamirian and M. Maul, Phys. Rev. D **61** (2000) 073004 [arXiv:hep-ph/9907495].
- [60] N. G. Stefanis, W. Schroers and H.-Ch. Kim, Eur. Phys. J. C **18** (2000) 137.
- [61] A. P. Bakulev, et al., Phys. Rev. D **70** (2004) 033014 [arXiv:hep-ph/0405062].
- [62] Z.-T. Wei and M.-Z. Yang, Phys. Rev. D **67** (2003) 094013.
- [63] T. Huang, X.-G. Wu and X.-H. Wu, Phys. Rev. D **70** (2004) 053007 [arXiv:hep-ph/0404163].
- [64] T. Huang and X.-G. Wu, Phys. Rev. D **70** (2004) 093013 [arXiv:hep-ph/0408252].
- [65] F. D. R. Bonnet et al. [Lattice Hadron Physics Collaboration], Phys. Rev. D **72** (2005) 054506 [arXiv:hep-lat/0411028].
- [66] S. Hashimoto et al. [JLQCD Collaboration], PoS **LAT2005**, (2006) 336 [arXiv:hep-lat/0510085].
- [67] D. Brömmel et al., PoS **LAT2005**, (2006) 360 [arXiv:hep-lat/0509133]; Eur. Phys. J. C **51** (2007) 335 [arXiv:hep-lat/0608021].
- [68] P.-H. J. Hsu and G. T. Fleming [arXiv:hep-lat/0710.4538].
- [69] A. P. Bakulev and A. V. Radyushkin, Phys. Lett. B **271** (1991) 223.
- [70] G. P. Lepage and S. J. Brodsky, Phys. Lett. B **87** (1979) 359; Phys. Rev. Lett. **43** (1979) 545; Phys. Rev. D **22** (1980) 2157; Perturbative Quantum Chromodynamics, A. H. Mueller ed., p.93, World Scientific, Singapore 1989; S. J. Brodsky, in Proceedings of the *Quantum Chromodynamics* Workshop, La Jolla, California, 1978.
- [71] A. V. Efremov, and A. V. Radyushkin, Phys. Lett. B **94** (1980) 245; Theor. Math. Phys. **42** (1980) 97.
- [72] A. V. Radyushkin, [arXiv:hep-ph/0410276].
- [73] A. Duncan and A. H. Mueller, Phys. Rev. D **21** (1980) 1636.
- [74] S. V. Mikhailov and A. V. Radyushkin, JETP Lett. **43** (1986) 712; Sov. J. Nucl. Phys. **49** (1989) 494; Phys. Rev. D **45** (1992) 1754.
- [75] A. P. Bakulev and A. V. Radyushkin, Phys. Lett. B **271** (1991) 223.
- [76] A. P. Bakulev and S. V. Mikhailov, Z. Phys. C **68** (1995) 451.
- [77] B. L. Ioffe and A. V. Smilga, Phys. Lett. B **114** (1982) 353.
- [78] N. Isgur and C. H. Llewellyn Smith, Phys. Rev. Lett. **52** (1984) 1080; Nucl. Phys. B **317** (1989) 526; Phys. Lett. B **217** (1989) 535.

- [79] A. V. Radyushkin, Acta. Phys. Polonica B **15** (1984) 403; Nucl. Phys. A **527** (1991) 153C; Nucl. Phys. A **532** (1991) 141.
- [80] O. C. Jacob and L. S. Kisslinger, Phys. Rev. Lett. **56** (1986) 225.
- [81] R. Jacob and P. Kroll, Phys. Lett. B **315** (1993) 463.
- [82] J. Bolz et al., Z. Phys. C **66** (1995) 267.
- [83] V. M. Braun and I. Halperin, Phys. Lett. B **328** (1994) 457.
- [84] B. Chibisov and A. R. Zhitnitsky, Phys. Rev. D **52** (1995) 5373.
- [85] D. Melikhov, Phys. Rev. D **53** (1996) 2460; Eur. Phys. J. direct C **4** (2002) 2 [arXiv:hep-ph/0110087].
- [86] S. Descotes and C. T. Sachrajda, Nucl. Phys. B **625** (2002) 239.
- [87] S. J. Brodsky, G. P. Lepage and T. Huang, Invited Talk at the Banff Summer Institute on Particle and Fields (1981), A. Z. Capri and A. N. Kamal, ed., p.143, Plenum Press, New York 1983.
- [88] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147** (1979) 385.
- [89] D. V. Shirkov and I. L. Solovtsov, Phys. Rev. Lett. **79** (1997), 1209; Phys. Lett. B **442** (1998), 344; Theor. Math. Phys. **120** (1999) 1220.
- [90] N. G. Stefanis, W. Schroers and H.-Ch. Kim, Phys. Lett. B **4429** (1999) 299. [arXiv:hep-ph/9807298].
- [91] H.-N. Li, Phys. Rev. D **52** (1995) 3958.
- [92] M. Beneke and Th. Feldmann, Nucl. Phys. B **592** (2001) 3.
- [93] Z. T. Wei and M. Z. Yang, Nucl. Phys. B **642** (2002) 263.
- [94] T. Kurimoto, H.-N. Li and A. I. Sanda, Phys. Rev. D **65** (2002) 014007.
- [95] H.-N. Li, Phys. Rev. D **66** (2002) 094010.